arXiv:1501.01094v2 [math.SG] 26 Jan 2015

Relative Invariants, Contact Geometry and Open String Invariants

An-Min Li[1]

Department of Mathematics, Sichuan University
Chengdu, PRC

Li Sheng[2]

Department of Mathematics, Sichuan University
Chengdu, PRC

Abstract

In this paper we propose a theory of contact invariants and open string invariants, which are generalizations of the relative invariants. We introduce two moduli spaces $\overline{\mathcal{M}}_A(M^+, C, g, m+\nu, \mathbf{y}, \mathbf{p}, (\mathbf{k}, \mathfrak{c}))$ and $\overline{\mathcal{M}}_A(M, L; g, m+\nu, \mathbf{y}, \mathbf{p}, \overrightarrow{\mu})$, prove the compactness of the moduli spaces and the existence of the invariants.

Contents

1 Introduction **2**

2 Symplectic manifolds with cylindrical ends **4**

2.1 Contact manifolds 4
2.2 Neighbourhoods of Lagrangian submanifolds 4
2.3 Cylindrical almost complex structures 6
2.4 J-holomorphic maps with finite energy 7
2.5 Periodic orbits of Bott-type 8

3 Weighted sobolev norms **11**

4 Moduli spaces of J-holomorphic maps **12**

4.1 Boundary conditions 12
4.2 Homology 13
4.3 Holomorphic blocks in M^+ 14
4.4 Holomorphic blocks in $\mathbb{R}\times\widetilde{M}$ 15
4.5 Local coordinate system of holomorphic blocks 17

5 Compactness theorems **17**

5.1 Bubble phenomenon 17
  5.1.1 Bound of the number of singular points 18
  5.1.2 Construction of the bubble tree for (a-2) 19
  5.1.3 Construction of bubble tree for (b) 21
  5.1.4 For the case of genus 0 22
5.2 $\mathcal{T}$- rescaling 23
5.3 Equivalent $\mathcal{DT}$- rescaling 26

---

[1] partially supported by a NSFC grant

[2] partially supported by a NSFC grant

5.4 Procedure of re-scaling 28
5.5 Weighted dual graph with a oriented decomposition 29
5.6 Li-Ruan's Compactification 32

6 Gluing theory-Pregluing **34**
6.1 Gluing one relative node for $M^+ \cup (\mathbb{R} \times \widetilde{M})$ 34
6.2 Gluing two relative nodal points for $M^+ \cup (\mathbb{R} \times \widetilde{M})$ 35
6.3 Norms on $C^\infty(\Sigma_{(r)}; u^*_{(r)}TM^+_{(r)})$ 37

7 Gluing theory–Regularization **38**
7.1 Local regularization 38
7.1.1 Top strata. 38
7.1.2 Lower strata. 39
7.2 Global regularization 41

8 Gluing theory– analysis estimates **42**
8.1 Gluing theory for 1-nodal case 42
8.2 Estimates of right inverse 42
8.3 Isomorphism between $KerD\mathcal{S}_b$ and $KerD\mathcal{S}_{b_{(r)}}$ 43
8.4 Gluing maps 46
8.5 Surjectivity and injectivity 47

9 Estimates of differentiations for gluing parameters **48**
9.1 Linear analysis on weight sobolov spaces 48
9.2 Estimates of $\frac{\partial Q_{b_{(r)}}}{\partial r}$ 49
9.3 Estimates of $\frac{\partial I_{(r)}}{\partial r}$ 51
9.4 Estimates of $\frac{\partial}{\partial r}\left[I_r(\kappa,\zeta) + Q_{b(r)}f_{(r)}(I_r(\kappa,\zeta))\right]$ 51

10 Contact invariants and Open string invariants **55**

1 Introduction

Open string invariant theory have been studied by many mathematicians and physicists (see [1,2, 11,16,17]). This theory closely relates to relative Gromov-Witten theory and contact geometry. In this paper we propose a theory of contact invariants and open string invariants, which are generalizations of the relative invariants. We outline the idea as follows.

1). Let (M,ω) be a compact symplectic manifold, $L \subset M$ be a compact Lagrangian submanifold. Let $(x_1,\cdots,x_n)$ be a local coordinate system in $O \subset L$, there is a canonical coordinates $(x_1,\cdots,x_n,y_1,\cdots,y_n)$ in $T^*L|_O$. Suppose that given a Riemannian metric g on L, and consider the unit sphere bundle $\widetilde{M}$. Let Λ be the Liouville form on T^*L, denote $\lambda = -\vartheta\mid_{\widetilde{M}}$. Then $(\widetilde{M},\lambda)$ is a contact manifold with contact form λ. The Reeb vector field is given by $X = \sum \frac{y_i}{\|y\|}\frac{\partial}{\partial x_i}$. By Lagrangian Neighborhood Theorem we can write $M - L$ as

$$M^+ = M_0 \bigcup \{[0,\infty) \times \widetilde{M}\},$$

where M_0 is a compact symplectic manifold with boundary. We choose an almost complex structure J on M^+ such that J is tamed by ω and over the cylinder end J is given by

$$J\mid_{\xi}=\tilde{J}, \quad JX=-\frac{\partial}{\partial a}, \;\; J(\frac{\partial}{\partial a})=X,$$

where $\xi=ker\lambda$, a is the canonical coordinate in $\mathbb{R}$, and $\tilde{J}$ is a $d\lambda$-tame almost complex structure in ξ. Assume that the periodic orbit sets of X are either non-degenerate or of Bott-type, and $\tilde{J}$ can be chosen such that $L_XJ=0$ along every periodic orbit.

Let $\overset{\circ}{\Sigma}$ be a Riemann surface with a puncture point p. We use the cylinder coordinates (s,t) near p, i.e.,we consider a neighborhood of p as $(s_0,\infty)\times S^1$. Let $u:\overset{\circ}{\Sigma}\rightarrow M^+$ be a J-holomorphic map with finite energy. Suppose that

$$[\lim_{s\rightarrow\infty}u(s,S^1)]=\sum\mu^i[c_i],$$

where $[c_i], i=1,...,\mathfrak{a}$, is a bases in $H_1(L,\mathbb{Z})$ and $\mu^i\in\mathbb{Z}$. Then $u(s,t)$ converges to a periodic orbit $x\subset\widetilde{M}$ of the Reeb vector field X as $s\rightarrow\infty$. We can view $\lim_{s\rightarrow\infty}u(s,S^1)$ as a loop in L, representing $\sum\mu^i[c_i]$. In this way, we can control the behaviour at infinity of J-holomorphic maps with finite energy.

2). To compactify the moduli space we need study the J-holomorphic maps into $\{(-\infty,\infty)\times\widetilde{M}\}$. There are two global vector fields on $\mathbb{R}\times\widetilde{M}$: $\frac{\partial}{\partial a}$ and X. Similar to the situation of relative invariants, there is a $\mathbb{R}$ action, which induces a $\mathbb{R}$-action on the moduli space of J-holomorphic maps. We need mod this action. Since there is a vector field X with $|X|=1$ on $\widetilde{M}$, the Reeb vector field, there is a one parameter group φ_θ action on $\widetilde{M}$ generated by $-X$. In particular, there is a S^1-action on every periodic orbit, corresponding to the freedom of the choosing origin of S^1. Along every periodic orbit we have $L_X\lambda=0$, on the other hand, we assume that on every periodic orbit $L_XJ=0$, then we can mod this action.

On the other hand, we choose the Li-Ruan's compactification in [18], that is, we firstly let the Riemann surfaces degenerate in Delingne-Mumford space and then let M^+ degenerate compatibly as in the situation of relative invariants. At any node, the Riemann surface degenerates independently with two parameters, which compatible with those freedoms of choosing the origins (see section §5 for degeneration and section §6 for gluing). Then both blowups at interior and at infinity lead boundaries of codimension 2 or more in the moduli space.

3). Another core technical issue in this paper is to define invariants using virtual techniques. As we know, there had been several different approaches, such as Fukaya-Ono [10], Li-Tian [21], Liu-Tian([22]), Ruan([27]), Siebert([29]) and etc. In [18], Li and Ruan provide a completely new approach to this issue: they show that the invariants can be defined via the integration on the *top* stratum virtually. In order to achieve this goal, they provide *refined* estimates of differentiations for gluing parameters: $\partial/\partial r$. In [18] the estimates for $\frac{\partial}{\partial r}$ is of order r^{-2} when $r\rightarrow\infty$, that is enough to define invariants. In this paper these estimates achieve to be of exponential decay order $\exp(-\mathfrak{c}r)$. Then we can use the estimates to define the invariants and prove the smoothness of the moduli space. The same method can be applied to GW-invariant for compact symplectic manifold.

In this paper we introduce two moduli spaces $\overline{\mathcal{M}}_A(M^+,C,g,m+\nu,\mathbf{y},\mathbf{p},(\mathbf{k},\mathfrak{e}))$ and $\overline{\mathcal{M}}_A(M,L;g,m+\nu,\mathbf{y},\mathbf{p},\overrightarrow{\mu})$, prove the compactness of the moduli spaces and the existence of the contact invariants

$\Psi^{(C)}_{(A,g,m+\nu,\mathbf{k},\mathfrak{c})}(\alpha_1,...,\alpha_m;\beta_{m+1},...,\beta_{m+\nu})$ and the open string invariants $\Psi^{(L)}_{(A,g,m+\nu,\overrightarrow{\mu})}(\alpha_1,...,\alpha_m)$. In our next paper [20] we will prove the smoothness of the two moduli spaces. Our open string invariants $\Psi^{(L)}_{(A,g,m+\nu,\overrightarrow{\mu})}$ can be generalized to L which is a disjoint union of compact Lagrangian sub-manifolds L_1,...,L_d.

We consider a neighborhood of Lagrangian sub-manifold L as $\mathbb{R}\times\widetilde{M}$. By the same method above we can define a local open string invariant. We will discuss this problem and calculate some examples in our next paper.

2 Symplectic manifolds with cylindrical ends

2.1 Contact manifolds

Let $(\mathcal{Q},\lambda)$ be a $(2n-1)$-dimensional compact manifold equipped with a contact form λ. We recall that a contact form λ is a 1-form on $\mathcal{Q}$ such that $\lambda\bigwedge(d\lambda)^{n-1}$ is a volume form. Associated to $(\mathcal{Q},\lambda)$ we have the contact structure $\xi=\ker(\lambda)$, which is a $(2n-2)$-dimensional subbundle of $T\mathcal{Q}$, and $(\xi,d\lambda|_\xi)$ defines a symplectic vector bundle. Furthermore, there is a unique nonvanishing vector field $X=X_\lambda$, called the Reeb vector field, defined by the condition

$$i_X\lambda=1,\quad i_Xd\lambda=0.$$

We have a canonical splitting of TQ,

$$T\mathcal{Q}=\mathbb{R}X\oplus\xi,$$

where $\mathbb{R}X$ is the line bundle generated by X.

2.2 Neighbourhoods of Lagrangian submanifolds

Let (M,ω) be a compact symplectic manifold, $L\subset M$ be a compact Lagrangian sub-manifold. The following Theorem is well-known.

Theorem 2.1. *Let (M,ω) be a symplectic manifold of dimension $2n$, and L be a compact Lagrangian submanifold. Then there exists a neighbourhood $U\subset T^*L$ of the zero section, a neighbourhood $V\subset M$ of L, and a diffeomorphism $\phi:U\to V$ such that*

$$\phi^*\omega=-d\Lambda,\quad \phi|_L=id, \tag{1}$$

where Λ is the canonical Liouville form.

Let $(x_1,\cdots,x_n)$ be a local coordinate system on $O\subset L$, there is a canonical coordinates

$$(x_1,\cdots,x_n,y_1,\cdots,y_n)$$

on $T^*O=T^*L|_O$. In terms of this coordinates the Liouville form can be written as

$$\Lambda=\sum y_idx_i.$$

Let $\pi:T^*L\to L$ be the canonical projection. There is a global defined vector field W in T^*L such that in the local coordinates of $\pi^{-1}(O)$, W can be written as

$$W|_{\pi^{-1}(O)}=-\sum_{i=1}^n y_i\frac{\partial}{\partial y_i}. \tag{2}$$

Suppose that given a Riemannian metric on L, in terms of the coordinates $x_1, ..., x_n$, $g_L = \sum\limits_{i,j=1}^{n} g_{ij} dx_i dx_j$. It naturally induced a metric on T^*L. Let

$$V = \frac{W}{\|W\|}.$$

V is a global defined vector field on $T^*L - L$.

Denote by $S^{n-1}(1)$ (resp.$B_1(0)$) the Euclidean unit sphere (resp. the Euclidean unit ball). Consider the coordinates transformation between the sphere coordinates and the Cartesian coordinate

$$\begin{aligned} \Psi : (0,1] \times S^{n-1}(1) &\to B_1(0) \\ (r, \theta_1, \cdots, \theta_{n-1}) &\to (y_1, \cdots, y_n). \end{aligned} \tag{3}$$

Consider the unit sphere bundle $\widetilde{M}$ and the unit ball bundle $\mathbb{D}_1(T^*L)$ in T^*L, in terms of the coordinates $(x_1, \cdots, x_n, y_1, \cdots, y_n)$

$$\widetilde{M}|_{\pi^{-1}(O)} = \{(x_1, \cdots, x_n, y_1, \cdots, y_n) \in \pi^{-1}(O) \mid \sum_{i,j=1}^{n} g^{ij}(x) y_i y_j = 1\}, \tag{4}$$

$$\mathbb{D}_1(T^*L)|_{\pi^{-1}(O)} = \{(x_1, \cdots, x_n, y_1, \cdots, y_n) \in \pi^{-1}(O) \mid \sum_{i,j=1}^{n} g^{ij}(x) y_i y_j \leq 1\}. \tag{5}$$

Denote $\lambda = -\Lambda \mid_{\widetilde{M}}$. We have

$$\Lambda = -\|y\|\lambda.$$

λ is a contact form, i.e., $(\widetilde{M}, \lambda)$ is a contact manifold. Put $\xi = \ker(\lambda)$. Then $X \mid_{\widetilde{M}} = -\sum g^{ij} y_i \frac{\partial}{\partial x_j}$ is the Reeb vector field, and $V \mid_{\widetilde{M}} = \sum_{i=1}^{n} y_i \frac{\partial}{\partial y_i}$.

The map Ψ induced a map $\tilde{\Psi} : (0,1] \times \widetilde{M} \to \mathbb{D}_1(T^*L)$. Through $\tilde{\Psi}$ we consider $\mathbb{D}_1(T^*L)|_{\pi^{-1}(O)} - L$ as $(0,1] \times \widetilde{M}$. By Theorem 2.1 we consider $M - L$ as

$$M^+ = M_0^+ \bigcup \{(0,1] \times \widetilde{M}\}$$

with the symplectic form

$$\omega_\phi = -d\Lambda = \|y\| d\lambda + d\|y\| \wedge \lambda, \tag{6}$$

where $M^+ := M - L$ and M_0^+ is a compact symplectic manifold with boundary.

We choose the *neck stretching technique.*

Let $\phi : [0, \infty) \to (0, \ell]$ be a smooth function satisfying, for any $k > 0$,

(1) $\phi' < 0, \;\; \phi(0) = \ell, \;\; \phi(a) \to 0 \;\; as \;\; a \to \infty,$

(2) $\lim\limits_{a \to 0^+} \frac{\partial^k \phi}{\partial a^k} = 0.$

Through ϕ we consider M^+ to be $M^+ = M_0^+ \bigcup \{[0, \infty) \times \widetilde{M}\}$ with symplectic form $\omega_\phi|_{M_0^+} = \omega$, and over the cylinder $[0, \infty) \times \widetilde{M}$

$$\omega_\phi = -d\Lambda = \phi d\lambda + \phi' da \wedge \lambda. \tag{7}$$

Moreover, if we choose the origin of $\mathbb{R}$ tending to ∞, we obtain $\mathbb{R} \times \widetilde{M}$ in the limit.

Choose $\ell_0 < \ell$ and denote

$$\Phi^+ = \left\{ \phi : [0, \infty) \to (0, \ell_0] | \phi' < 0 \right\}.$$

Let $\ell_1 < \ell_2$ be two real numbers satisfying $0 < \ell_1 < \ell_2 \leq \ell_0$. Let Φ_{ℓ_1,ℓ_2} be the set of all smooth functions $\phi : \mathbb{R} \to (\ell_1, \ell_2)$ satisfying

$$\phi' < 0, \quad \phi(a) \to \ell_1 \quad \text{as } a \to \infty, \quad \phi(a) \to \ell_2 \quad \text{as } a \to -\infty.$$

To simplify notations we use Φ to denote both Φ^+ and Φ_{ℓ_1,ℓ_2}, in case this does not cause confusion.

We fixed $\phi \in \Phi$ and consider the symplectic manifold $\left(M_0^+ \bigcup \{[0, \infty) \times \widetilde{M}\}, \omega_\phi \right)$. For any different $\phi_1 \in \Phi$, we have $\omega_\phi|_{M_0^+} = \omega_{\phi_1}|_{M_0^+}$ and $\phi \circ \phi_1^{-1}$ is a symplectic diffeomorphic over cylinder part $\{(0, 1] \times \widetilde{M}\}$.

2.3 Cylindrical almost complex structures

Let

$$M^+ = M_0^+ \bigcup \left\{ [0, \infty) \times \widetilde{M} \right\} \tag{8}$$

be a symplectic manifold with cylindrical end, where $\widetilde{M}$ be a compact contact manifold with contact form λ. Denote by ω_ϕ the symplectic form of M^+ such that $\omega_\phi|_{M_0^+} = \omega$, and over the cylinder $[0, \infty) \times \widetilde{M}$

$$\omega_\phi = -d\Lambda = \phi d\lambda + \phi' da \wedge \lambda. \tag{9}$$

We also consider $\mathbb{R} \times \widetilde{M}$. Denote by N one of M^+ and $\mathbb{R} \times \widetilde{M}$.

Put $\xi = \ker(\lambda)$, and denote by X the Reeb vector field defined by

$$\lambda(X) = 1, \quad d\lambda(X) = 0.$$

We choose a $d\lambda$-tame almost complex structure $\widetilde{J}$ for the symplectic vector bundle $(\xi, d\lambda) \to \widetilde{M}$ such that

$$g_{\widetilde{J}(x)}(h, k) = \frac{1}{2} \left(d\lambda(x)(h, \widetilde{J}(x)k) + d\lambda(x)(k, \widetilde{J}(x)h) \right), \tag{10}$$

for all $x \in \widetilde{M},\ h, k \in \xi_x$, defines a smooth fibrewise metric for ξ. We assume that we can choose $\widetilde{J}$ such that on every periodic orbit $L_X \widetilde{J} = 0$. Denote by $\Pi : T\widetilde{M} \to \xi$ the projection along X. We define a Riemannian metric $\langle\ ,\ \rangle$ on $\widetilde{M}$ by

$$\langle h, k \rangle = \lambda(h)\lambda(k) + g_{\widetilde{J}}(\Pi h, \Pi k) \tag{11}$$

for all $h, k \in T\widetilde{M}$.

Given a $\widetilde{J}$ as above there is an associated almost complex structure J on $\mathbb{R} \times \widetilde{M}$ defined by

$$J \mid_\xi = \tilde{J}, \quad JX = \frac{\partial}{\partial a}, \quad J(\frac{\partial}{\partial a}) = -X, \tag{12}$$

where a is the canonical coordinate in $\mathbb{R}$. It is easy to check that J defined by (12) is ω_φ-tame over the cylinder end. We can choose an almost complex structure J on M^+ such that J is tamed by ω and over the cylinder end J is given by (12).

There is a canonical coordinate system for $\mathbb{R} \times \widetilde{M}$ and for cylinder end of M^+, but still there are some freedom of choosing the coordinates:

When we write M^+ as (8) we have chosen a coordinate a over the cylinder part. We can choose different coordinate $\hat{a}$ over the cylinder part such that

$$a = \hat{a} + C \tag{13}$$

for some constant $C > 0$. Similarly, for $\mathbb{R} \times \widetilde{M}$ we can choose $\hat{a}$ such that

$$a = \hat{a} \pm C \tag{14}$$

for some constant $C > 0$.

For any $\phi \in \Phi$

$$\langle v, w \rangle_{\omega_\phi} = \frac{1}{2} \left(\omega_\phi(v, Jw) + \omega_\phi(w, Jv) \right) \qquad \forall \;\; v, w \in TN \tag{15}$$

defines a Riemannian metric on N. Note that $\langle \, , \, \rangle_{\omega_\phi}$ is not complete. We choose another metric $\langle \, , \, \rangle$ on N such that

$$\langle \, , \, \rangle = \langle \, , \, \rangle_{\omega_\phi} \quad \textit{on} \;\; M_0^+ \tag{16}$$

and over the tubes

$$\langle (a, v), (b, w) \rangle = ab + \lambda(v)\lambda(w) + g_{\widetilde{J}}(\Pi v, \Pi w), \tag{17}$$

where we denote by $\Pi : T\widetilde{M} \to \xi$ the projection along X. It is easy to see that $\langle \, , \, \rangle$ is a complete metric on N.

2.4 J-holomorphic maps with finite energy

Let (Σ, i) be a compact Riemann surface and $P \subset \Sigma$ be a finite collection of puncture points. Denote $\overset{\circ}{\Sigma} = \Sigma \backslash P$. Let $u : \overset{\circ}{\Sigma} \to N$ be a J-holomorphic map, i.e., u satisfies

$$du \circ i = J \circ du. \tag{18}$$

Following [14] we impose an energy condition on u. For any J-holomorphic map $u : \overset{\circ}{\Sigma} \to N$ and any $\phi \in \Phi$ the energy $E_\phi(u)$ is defined by

$$E_\phi(u) = \int_\Sigma u^* \omega_\phi. \tag{19}$$

Let $z = e^{s + 2\pi\sqrt{-1}t}$. One computes over the cylindrical part

$$u^*\omega_\phi = \left(\phi d\lambda \left((\pi\widetilde{u})_s, (\pi\widetilde{u})_t \right) - \phi'(a_s^2 + a_t^2) \right) ds \wedge dt, \tag{20}$$

which is a nonnegative integrand. A J-holomorphic map $u : \overset{\circ}{\Sigma} \to N$ is called a finite energy J-holomorphic map if over the cylinder end

$$\sup_{\phi \in \Phi} \left\{ \int_\Sigma u^* \omega_\phi \right\} + \int_\Sigma u^* d\lambda < \infty. \tag{21}$$

For a J-holomorphic map $u : \Sigma \to \mathbb{R} \times \widetilde{M}$ we write $u = (a, \widetilde{u})$ and define

$$\widetilde{E}(u) = \int_{\Sigma} \widetilde{u}^* d\lambda. \tag{22}$$

Denote

$$\widetilde{E}(s) = \int_s^{\infty} \int_{S^1} \widetilde{u}^*(d\lambda).$$

Then

$$\widetilde{E}(s) = \int_s^{\infty} \int_{S^1} |\Pi \widetilde{u}_t|^2 ds dt,$$

$$\frac{d\widetilde{E}(s)}{ds} = -\int_{S^1} |\Pi \widetilde{u}_t|^2 dt. \tag{23}$$

Here and later we use $|\cdot|$ denotes the norm with respect to the metric defined by (17).

The following two lemmas are well-known (see [13]):

Lemma 2.2. *(1) Let $u = (a, \widetilde{u}) : \mathbb{C} \to \mathbb{R} \times \widetilde{M}$ be a J-holomorphic map with finite energy. If $\int_{\mathbb{C}} \widetilde{u}^*(\pi^* d\lambda) = 0$, then u is a constant.*

(2) Let $u = (a, \widetilde{u}) : \mathbb{R} \times S^1 \to \mathbb{R} \times \tilde{M}$ be a J-holomorphic map with finite energy. If $\int_{\mathbb{R} \times S^1} \widetilde{u}^*(\pi^* d\lambda) = 0$, then $(a, \widetilde{u}) = (kTs + c, kt + d)$, where $k \in \mathbb{Z}^+$, c and d are constants.

Lemma 2.3. *Let $u = (a, \widetilde{u}) : \mathbb{C} - D_1 \to \mathbb{R} \times \widetilde{M}$ be a nonconstant J-holomorphic map with finite energy. Put $z = e^{s + 2\pi\sqrt{-1}t}$. Then for any sequence $s_i \to \infty$, there is a subsequence, still denoted by s_i, such that*

$$\lim_{i \to \infty} \widetilde{u}(s_i, t) = x(kTt)$$

in $C^{\infty}(S^1)$ for some kT-periodic orbit $x(kTt)$.

2.5 Periodic orbits of Bott-type

Let $\mathcal{F} \subset \widetilde{M}$ be the locus of minimal periodic orbits with $\mathcal{F} = \bigcup_{i=1}^{\ell} \mathcal{F}_i$, where each $\mathcal{F}_i$ is a connected component of $\mathcal{F}$ with minimal periodic T_i. We assume that

Assumption 2.4. *(1) every $\mathcal{F}_i$ is either non-degenerate or of Bott-type;*

(2) let $\mathcal{F}_i$ be of Bott-type, then there exists a free S^1-action on $\mathcal{F}_i$ such that $Z_i = \mathcal{F}_i / S^1$ is a closed, smooth manifold. Set $n_i = \dim(\mathcal{F}_i)$;

(3) for every periodic orbit x, there is a smooth submanifold $\Re_x \subset \widetilde{M}$ of dimension > 2 such that $d\lambda \mid_{\Re_x} = 0$ and $x \subset \Re_x$, and the almost complex structure $\tilde{J}$ can be chosen such that $L_X \tilde{J} = 0$ along x.

Let $u = (a, \widetilde{u}) : \mathbb{C} - D_1 \to \mathbb{R} \times \widetilde{M}$ be a J-holomorphic map with finite energy. Put $z = e^{s + 2\pi i t}$. Assume that there exists a sequence $s_i \to \infty$ such that $\tilde{u}(s_i, t) \longrightarrow x(kTt)$ in $C^{\infty}(S^1, \widetilde{M})$ as $i \to \infty$ for some $k \in \mathbb{Z}$, where T is the minimal periodic. Following Hofer (see [14]) we introduce a convenient local coordinates near the periodic orbit x. Since $S^1 = \mathbb{R}/\mathbb{Z}$, we work in the covering space $\mathbb{R} \to S^1$.

Lemma 2.5. *Let $(\widetilde{M},\lambda)$ be a $(2n-1)$-dimensional compact manifold, and let $x(kt)$ be a k-periodic orbit with the minimal periodic T. Then there is an open neighborhood $U\subset S^1\times\mathbb{R}^{2n-2}$ of $S^1\times\{0\}$ with coordinates $(\vartheta,w_1,\cdots,w_{2n-2})$ and an open neighborhood $\Im\subset\widetilde{M}$ of $\{x(t)|t\in\mathbb{R}\}$ and a diffeomorphism $\psi:U\to\Im$ mapping $S^1\times\{0\}$ onto $\{x(t)|t\in\mathbb{R}\}$ such that*

$$\psi^*\lambda=g\lambda_0, \tag{24}$$

where $\lambda_0=d\vartheta+\sum w_i dw_{n+i-1}$ and $g:U\to\mathbb{R}$ is a smooth function satisfying

$$g(\vartheta,0)=T,\ \ dg(\vartheta,0)=0 \tag{25}$$

for all $\vartheta\in S^1$.

Remark 2.6. *We call the coordinate system $(\vartheta,\mathbf{w})$ in Lemma 2.5 a pseudo-Darboux coordinate system, and call the following transformation of two local pseudo-Darboux coordinate systems*

$$(\vartheta,\mathbf{w})\longrightarrow(\check{\vartheta},\mathbf{w}),\ \ \check{\vartheta}=\vartheta+\vartheta_0 \tag{26}$$

a canonical coordinate transformation, where ϑ_0 is a constant.

The following theorem is well-known (see [5, 18, 32])

Theorem 2.7. *Suppose that $\mathcal{F}$ satisfies Assumption 2.4. Let $u:\mathbb{C}-D_1\to\mathbb{R}\times\tilde{M}$ be a J-holomorphic map with finite energy. Put $z=e^{s+2\pi\sqrt{-1}t}$. Then*

$$\lim_{s\to\infty}\widetilde{u}(s,t)=x(kTt)$$

in $C^\infty(S^1)$ for some kT-periodic orbit x, and there are constants ℓ_0, ϑ_0 such that for any $0<\mathfrak{c}<\min\{\frac{1}{2},\frac{\mathcal{C}_1^2}{2}\}$ and for all $\mathbf{n}=(n_1,n_2)\in\mathbb{Z}^2_{\geq 0}$

$$|\partial^{\mathbf{n}}[a(s,t)-kTs-\ell_0]|\leq\mathcal{C}_{\mathbf{n}}e^{-\mathfrak{c}|s|} \tag{27}$$

$$|\partial^{\mathbf{n}}[\vartheta(s,t)-kt-\vartheta_0]|\leq\mathcal{C}_{\mathbf{n}}e^{-\mathfrak{c}|s|} \tag{28}$$

$$|\partial^{\mathbf{n}}\mathbf{w}(s,t)|\leq\mathcal{C}_{\mathbf{n}}e^{-\mathfrak{c}|s|}, \tag{29}$$

where $C_{\mathbf{n}}$ are constants. Here $(\vartheta,\mathbf{w})$ is a pseudo-Darboux coordinate near the periodic orbit x.

Let x_o be a minimal periodic orbit. We choose a local pseudo-Darboux coordinate on an open set $\Im\subset\widetilde{M}$ near x_o. Let $\mathcal{O}$ be an open set such that $\overline{\mathcal{O}}\subset\Im$ is compact and $x_o\subset\mathcal{O}$. We fix a positive constant C. Denote by $\mathfrak{S}$ the class of J-holomorphic maps with finite energy $u:[s_0',\infty)\times S^1\to\mathbb{R}\times\tilde{M}$ satisfying

(i) $\sup_{\phi\in\Phi}E_\phi(u)\leq\frac{1}{2}\hbar$, where $\hbar$ is the constant in Theorem 5.2 and Lemma 5.3,

(ii) $\lim_{s\to\infty}\widetilde{u}(s,t)=x(kTt)$ in $C^\infty(S^1)$ with $x\subset\mathcal{O}$,

(iii) $\widetilde{u}([s_0',\infty)\times S^1)$ lie in the pseudo-Darboux coordinate system on $\Im$,

(iv) there is a ball $D_1(s_0,t_0)\subset[s_0',\infty)\times S^1$ such that $\widetilde{u}(D_1(s_0,t_0))\subset\mathcal{O}$ and $|a(D_1(s_0,t_0))|\leq\mathsf{C}$.

In our next paper [19] we will prove the following theorem.

Theorem 2.8. *Let $u \in \mathfrak{S}$. Then the constants $C_{\mathbf{n}}$ in Theorem 2.7 depend only on $\mathcal{C}_1, s_0, \mathsf{C}$, $\mathbf{n}$, $\hbar$, $\mathfrak{c}$ and $\mathcal{O}$. Moreover, we have*

$$|\ell_0| \le C_1, \tag{30}$$

where C_1 depends only on $\hbar$, $C_{\mathbf{n}}, \mathfrak{c}, s_0, \mathsf{C}$ and $\mathcal{O}$.

Following [14] we introduce functions

$$a^{\diamond}(s,t) = a(s,t) - ks, \quad \vartheta^{\diamond}(s,t) = \vartheta(s,t) - kt. \tag{31}$$

Denote

$$\pounds = (a^{\diamond}, \vartheta^{\diamond}). \tag{32}$$

Set $b_i = 0,\ b_{n-1+i}(\mathbf{w}) = w_i, \forall\ i = 1, \cdots, n-1$, and

$$e_i = \frac{\partial}{\partial w_i} - b_i \frac{\partial}{\partial \vartheta}, \quad i = 1, \cdots, 2n-2.$$

Then $\xi =$span$\{e_1, \cdots, e_{2n-2}\}$. Denote $\tilde{J} e_i = \sum \tilde{J}_{ij} e_i$. In the basis $\partial_\vartheta, e_1, \cdots, e_{2n-2}$, the Reeb vector field can be re-written as

$$X = \frac{1}{g}\partial_\vartheta + \frac{1}{g^2}\left(\sum_{i \le n-1} e_{n-1+i}(g) e_i - \sum_{i \ge n} e_{i-n+1}(g) e_i\right). \tag{33}$$

Let $\tilde{\mathbf{X}} = \frac{1}{g^2}(e_n g, \cdots, e_{2n-2} g, -e_1 g, \cdots, -e_{n-1} g)$. By (18), we have

$$\pounds_s + J \pounds_t = h, \quad \mathbf{w}_s + \widetilde{J} \mathbf{w}_t + a_t \tilde{\mathbf{X}} - a_s \tilde{J} \tilde{\mathbf{X}} = 0.$$

where

$$h = (\Sigma b_i(\mathbf{w})(w_i)_t + (\vartheta_t + \Sigma b_i(\mathbf{w})(w_i)_t)(\Sigma f_i w_i), -\Sigma b_i(\mathbf{w})(w_i)_s - (\vartheta_t + \Sigma b_i(\mathbf{w})(w_i)_s)(\Sigma f_i w_i))$$

and $f_i = \int_0^1 \partial_{w_i} g(\vartheta, \tau \mathbf{w}) d\tau$. Let $V = \pounds_t$ and $g = h_t$. Denote

$$\hat{E}(\pounds) = \int_\Sigma \|\pounds_s\|^2 + \|\pounds_t\|^2 ds dt. \tag{34}$$

In our next paper [19] we will prove the following theorem.

Theorem 2.9. *Suppose that $\widetilde{M}$ satisfies the Assumption 2.4. Let $u : [-R, R] \times S^1 \to \mathbb{R} \times \widetilde{M}$ be a J-holomorphic maps with finite energy. Assume that*

(i) $\sup\limits_{\phi \in \Phi} E_\phi(u, -R \le s \le R) + \hat{E}(\pounds) \le \frac{1}{2}\hbar$,

(ii) $\widetilde{u}([-R, R] \times S^1)$ *lie in a pseudo-Darboux coordinate system* $(\vartheta, \mathbf{w})$ *on* $\Im$,

(iii) $\sum\limits_{n_1, n_2 \le 2} \|\nabla^{\mathbf{n}} u(-R, \cdot)\|_{L^2(S^1)} \le C_2, \qquad \sum\limits_{n_1, n_2 \le 2} \|\nabla^{\mathbf{n}} u(R, \cdot)\|_{L^2(S^1)} \le C_2$, *where* $\mathbf{n} = (n_1, n_2)$,

Then there exist constants $\mathcal{C}_1 > 0$ and $0 < \mathfrak{c} < \frac{1}{2}$ depending only on $\tilde{J}$ and C_2 such that

$$|\nabla \mathbf{w}|(s,t) \le \mathcal{C}_1 e^{-\mathfrak{c}(R - |s|)}, \quad |\nabla \pounds| \le \mathcal{C}_1 e^{-\mathfrak{c}(R - |s|)}, \quad \forall\ |s| \le R - 1, \tag{35}$$

In this paper, we also need the following implicit function theorem (see [23]).

Lemma 2.10. *Let X and Y be Banach spaces, $U \subset X$ be an open set, and ℓ be a positive integer. If $F : U \longrightarrow Y$ is of class C^{ℓ}. Let $x_0 \in U$ be such that $D := dF(x_0) : X \longrightarrow Y$ is surjective and has a bounded linear right inverse $Q : Y \longrightarrow X$. Choose positive constants $\hbar_1$ and C such that $\|Q\| \leq C$, $B_{\hbar_1}(x_0, X) \subset U$ and*

$$\|dF(x) - D\| \leq \frac{1}{2C}, \quad \forall\, x \in B_{\hbar_1}(x_0) \tag{36}$$

where $B_{\hbar_1}(x_0) = \{x \in X |\ \|x - x_0\| \leq \hbar_1\}$. Suppose that $x_1 \in X$ satisfies

$$\|F(x_1)\| < \frac{\hbar_1}{4C}, \quad \|x_1 - x_0\| \leq \frac{\hbar_1}{8}. \tag{37}$$

Then there exists a unique $x \in X$ such that

$$F(x) = 0, \quad x - x_1 \in Im\, Q, \quad \|x - x_0\| \leq \hbar_1, \quad \|x - x_1\| \leq 2C\|F(x_1)\|. \tag{38}$$

Moreover, write $x := x_0 + \xi + Q \circ f(\xi)$, $\xi \in kerD$, then f is of class C^{ℓ}.

3 Weighted sobolev norms

Consider $\mathbb{R} \times \widetilde{M}$ and $M^+ = M_0^+ \bigcup \left\{[0, \infty) \times \widetilde{M}\right\}$. Let N be one of $\mathbb{R} \times \widetilde{M}$, M^+. Suppose that $\Sigma = \bigcup\limits_v \Sigma_v$ is Riemann surface with nodal points $\{q_1, \cdots, q_{\mathfrak{j}}\}$, puncture points $\{p_1, \cdots, p_\nu\}$ and $u : \Sigma \to \bigcup N_i$ is a continuous map such that the restriction of u to each smooth component is smooth, where $\bigcup N_i$ denotes the union of some copy of N. We choose cylinder coordinates (s, t) on Σ near each nodal point and each puncture point. We choose a local pseudo-Darboux coordinate system near each periodic orbit on N. Let $\overset{\circ}{\Sigma} = \Sigma - \{q_1, \cdots, q_{\mathfrak{j}}, p_1, \cdots, p_\nu\}$.

Over each tube the linearized operator D_u takes the following form

$$D_u = \frac{\partial}{\partial s} + J_0 \frac{\partial}{\partial t} + S = \bar{\partial}_J + S. \tag{39}$$

By exponential decay we have

$$\left| \frac{\partial^{k+l}}{\partial s^k \partial t^l} S \right| \leq C_{k,l} e^{-\mathfrak{c}s}$$

for some constant $C_{k,l} > 0$ for s big enough. Therefore, the operator $H_s = J_0 \frac{d}{dt} + S$ converges to $H_\infty = J_0 \frac{d}{dt}$. Obviously, the operator D_u is not Fredholm operator because over each puncture and node the operator $H_\infty = J \frac{D}{dt}$ has zero eigenvalue. The $\ker H_\infty$ consists of constant vectors. To recover a Fredholm theory we use weighted function spaces. We choose a weight α for each end. Fix a positive function W on Σ which has order equal to $e^{\alpha|s|}$ on each end, where α is a small constant such that $0 < \alpha < \mathfrak{c}$ and over each end $H_\infty - \alpha = J_0 \frac{d}{dt} - \alpha$ is invertible. We will write the weight function simply as $e^{\alpha|s|}$. Denote by $C(\Sigma; u^*T(\bigcup N_i))$ all tangent vector fields h on $\bigcup N_i$ along u satisfying

(a) $h \in C^0(\Sigma, u^*T(\bigcup N_i))$,

(b) the restriction of h to each smooth component is smooth.

For any section $h \in C(\Sigma; u^*T(\bigcup N_i))$ and section $\eta \in \Omega^{0,1}(u^*T(\bigcup N_i))$ we define the norms

$$\|h\|_{1,p,\alpha} = \sum_v \left(\int_{\Sigma_v} (|h|^p + |\nabla h|^p) d\mu \right)^{1/p} + \sum_v \left(\int_{\Sigma_v} e^{2\alpha|s|} (|h|^2 + |\nabla h|^2) d\mu \right)^{1/2} \tag{40}$$

$$\|\eta\|_{p,\alpha} = \sum_v \left(\int_{\Sigma_v} |\eta|^p d\mu \right)^{1/p} + \sum_v \left(\int_{\Sigma_v} e^{2\alpha|s|} |\eta|^2 d\mu \right)^{1/2} \tag{41}$$

for $p \geq 2$, where all norms and covariant derivatives are taken with respect to the metric $\langle \ \rangle$ on $u^*T(\bigcup N_i)$ defined in (17), and the metric on $\overset{\circ}{\Sigma}$. Denote

$$\mathcal{C}(\Sigma; u^*T(\bigcup N_i)) = \{h \in C(\Sigma; u^*T(\bigcup N_i)); \|h\|_{1,p,\alpha} < \infty\}, \tag{42}$$

$$\mathcal{C}(u^*T(\bigcup N_i) \otimes \wedge^{0,1}) = \{\eta \in \Omega^{0,1}(u^*T(\bigcup N_i)); \|\eta\|_{p,\alpha} < \infty\}. \tag{43}$$

Denote by $W^{1,p,\alpha}(\Sigma; u^*T(\bigcup N_i))$ and $L^{p,\alpha}(u^*T(\bigcup N_i) \otimes \wedge^{0,1})$ the completions of $\mathcal{C}(\Sigma; u^*T(\bigcup N_i))$ and $\mathcal{C}(u^*T(\bigcup N_i) \otimes \wedge^{0,1})$ with respect to the norms (40) and (41) respectively. Then the operator $D_u : W^{1,p,\alpha} \to L^{p,\alpha}$ is a Fredholm operator.

For each puncture point $p_j, j = 1, ..., \nu$, let $h_{j0} \in (T_{p_j}(\mathcal{F}_{\mathfrak{e}_j}) \oplus (span\{\frac{\partial}{\partial a}\})$. For each bounded nodal q_i, denote $\mathbb{H}_{q_i} = T_{q_i}N$, let $h_{(i+\nu)0} \in \mathbb{H}_{q_i}$; for each unbounded nodal q_i, denote $\mathbb{H}_{q_i} = (T_{q_i}(\mathcal{F}_{i_j}) \oplus (span\{\frac{\partial}{\partial a}\})$ and let $h_{(i+\nu)0} \in \mathbb{H}_{q_i}$, where $u : \Sigma \to N$ converges to kT periodic orbit $x(kTt) \subset \mathcal{F}_{i_j}$ as $z \to q_i$. Put $\mathbb{H} = \left(\oplus_{j=1}^{\nu}(T_{p_j}(\mathcal{F}_{\mathfrak{e}_j}) \oplus (span\{\frac{\partial}{\partial a}\})\right) \bigoplus \left(\oplus_{i=1}^{\mathfrak{I}} \mathbb{H}_{q_i})\right)$, $h_0 = (h_{10}, ..., h_{\nu 0}, h_{(1+\nu)0}, ..., h_{(\mathfrak{I}+\nu)0})$.

h_0 may be considered as a vector field in the coordinate neighborhood. We fix a cutoff function ρ:

$$\rho(s) = \begin{cases} 1, & if\ |s| \geq d, \\ 0, & if\ |s| \leq \frac{d}{2} \end{cases}$$

where d is a large positive number. Put

$$\hat{h}_0 = \rho h_0.$$

Then for d big enough $\hat{h}_0$ is a section in $C^\infty(\Sigma; u^*TN)$ supported in the tube $\{(s,t) | |s| \geq \frac{d}{2}, t \in S^1\}$. Denote

$$\mathcal{W}^{1,p,\alpha} = \{h + \hat{h}_0 | h \in W^{1,p,\alpha}, h_0 \in \mathbb{H}\}.$$

We define the weighted Sobolev norm on $\mathcal{W}^{1,p,\alpha}$ by

$$\|(h, \hat{h}_0)\|_{\Sigma,1,p,\alpha} = \|h\|_{\Sigma,1,p,\alpha} + |h_0|.$$

Obviously, the operator $D_u : \mathcal{W}^{1,p,\alpha} \to L^{p,\alpha}$ is also a Fredholm operator.

4 Moduli spaces of J-holomorphic maps

4.1 Boundary conditions

Consider the symplectic manifold with cylindrical end

$$M^+ = M_0^+ \bigcup \left\{ [0, \infty) \times \widetilde{M} \right\}.$$

Let $((\Sigma, \mathbf{j}); \mathbf{y}, \mathbf{p})$ be a connected semistable curve with m marked points $\mathbf{y} = (y_1, ..., y_m)$ and ν puncture points $\mathbf{p} = (p_1, ..., p_\nu)$, and $u : \Sigma \to M^+$ be a J-holomorphic map. Let $\Sigma = \bigcup_{v=1}^{d} (\Sigma_v, j_v)$ where (Σ_v, j_v) is a smooth Riemann surface and $\pi_v : \Sigma_v \to \Sigma$ is a continuous map. To describe the boundary conditions we consider two different cases separately:

Case A . Moduli space of J-holomorphic maps in contact geometry.

Definition 4.1. *Let $\mathbf{p} = (p_1, \cdots, p_\nu)$ be the puncture points. We assign two weights $(\mathbf{k}, \mathfrak{e})$ to $\mathbf{p}$:*

(1) $\mathbf{k} : \mathbf{p} \to \mathbb{Z}_{>0}$ assigning a k_i to each puncture point p_i, denote $\mathbf{k} = (k_1, \cdots, k_\nu)$.

(2) $\mathfrak{e} : \mathbf{p} \to \mathbb{Z}_{>0}$ assigning a number $\mathfrak{e}_{\mathfrak{i}}$, $1 \leq \mathfrak{e}_{\mathfrak{i}} \leq \ell$ to each puncture point p_i.

We call a J-holomorphic map u satisfies $(\mathbf{k}, \mathfrak{e})$ boundary condition if

$u(z)$ converges to a $k_i \cdot T_{\mathfrak{e}_i}$-periodic orbit $x(k_i \cdot T_{\mathfrak{e}_i} t) \subset \mathcal{F}_{\mathfrak{e}_{\mathfrak{i}}}$ as z tends to p_i.

Case B. Moduli space of J-holomorphic maps in (M, L).

As we show in section §2.2 that $M - L$ can be considered as $M^+ = M_0^+ \bigcup \left\{ [0, \infty) \times \widetilde{M} \right\}$. Let $[c_i], i = 1, \cdots, \mathfrak{a}$ is a bases in $H_1(L; \mathbb{Z})$.

Definition 4.2. *Let $\mathbf{p} = (p_1, \cdots, p_\nu)$ be the order puncture points. We assign a weight $\overrightarrow{\mu}$ to $\mathbf{p}$: $\overrightarrow{\mu} : \mathbf{p} \to \mathbb{Z}_{>0}^{\oplus \mathfrak{a}}$ assigning a $\mu_i = \sum_{l=1}^{\mathfrak{a}} \mu_{il} [c_l]$ to each puncture point p_i, where $\mu_{il} \in \mathbb{Z}$. Choose the cylinder coordinates (s_i, t_i) near p_i. We call a J-holomorphic map u satisfies $(\overrightarrow{\mu})$ boundary condition if u satisfies*

$$[\pi(\lim_{s_i \to \infty} u(s_i, S^1))] = \mu_i, \ \ \forall\, 1 \leq i \leq \nu, \tag{44}$$

where $\pi : T^*L \to L$ is the canonical projection.

4.2 Homology

We fix $A \in H^2(M^+, \Re; \mathbb{Z})$ satisfying $\partial A = \sum [x(k_i T_{\mathfrak{e}_i} t)]$, where $x(k_i T_{\mathfrak{e}_i} t) \subset \mathcal{F}_{\mathfrak{e}_i}$ is a $k_i T_{\mathfrak{e}_i}$ periodic orbit. Consider a J-holomorphic map u satisfying

$$[u_*(\Sigma)] = A. \tag{45}$$

We show that the homology class A give a bound of Energy. To simplify notation we let $(u, (\Sigma, \mathbf{j}), \mathbf{y}, p)$ be a J-holomorphic map converging to a kT-periodic orbit $x(kTt)$ as z tends to p, where $x(kTt)$ lie in $\mathcal{F}_i$ and T is the minimal periodic. For the case **A**, by the assumption (3) of $\mathcal{F}$, we construct a connected surface $W \subset \Re_x$ with boundary $x(kTt)$ (including the kT-periodic). Then $u_*(\Sigma) \cup W$ is a closed surface in M^+ and

$$[u_*(\Sigma) \cup W] \in H^2(M^+; \mathbb{Z}).$$

Denote $A = [u_*(\Sigma) \cup W]$. By $d\lambda|_{\Re_x} = 0$ and $W \subset \widetilde{M}$ we have

$$\omega(A) = \int_{u_*(\Sigma)} \omega + \int_W \omega = E_\phi(u) + \int_W d\lambda = E_\phi(u). \tag{46}$$

Let $W' \subset \Re_x$ be another surface with boundary x, denote $A' = [u_*(\Sigma) \cup W'] \in H^2(M^+; \mathbb{Z})$. We have $\omega(A) = \omega(A') = E_\phi(u)$, that is, $E_\phi(u)$ is independent of the choice of W in $\Re_x$.

For case **B** let $A \in H^2(M, L; \mathbb{Z})$ be a fixed homology class satisfying $\partial A = \sum \mu_i$. We have the same results.

4.3 Holomorphic blocks in M^+

Let $((\Sigma,\mathbf{j});\mathbf{y},\mathbf{p})$ be a connected semistable curve with m marked points $\mathbf{y}=(y_1,...,y_m)$ and ν puncture points $\mathbf{p}=(p_1,...,p_\nu)$. Let $u:\Sigma\to M^+$ be a J-holomorphic map. Suppose that $u(z)$ converges to a $k_i\cdot T_{\mathfrak{e}_i}$-periodic orbit $x(k_iT_{\mathfrak{e}_i}t)\subset\mathcal{F}_{\mathfrak{e}_i}$ as z tends to p_i.

Definition 4.3. *A J-holomorphic map $(u;((\Sigma,\mathbf{j}),\mathbf{y},\mathbf{p}))$ is said to be stable if for each v one of the following conditions holds:*

(1). $u\circ\pi_{\Sigma_v}:\Sigma_v\to M^+$ is not a constant map.

(2). Let val_v be the number of special points on Σ_v which are nodal points, marked points or puncture points. Then $val_v+2g_v\geq 3$.

Definition 4.4. *Two stable J-holomorphic maps $\Gamma=(u,(\Sigma,\mathbf{j}),\mathbf{y},\mathbf{p})$ and $\check{\Gamma}=(\check{u},(\check{\Sigma},\check{\mathbf{j}}),\check{\mathbf{y}},\check{\mathbf{p}})$ is called equivalent if there exists a diffeomorphism $\varphi:\Sigma\to\check{\Sigma}$ such that it can be lifted to bi-holomorphic isomorphisms $\varphi_{vw}:(\Sigma_v,j_v)\to(\check{\Sigma}_w,\check{j}_w)$ for each component Σ_v of Σ, and*

(1) *$\varphi(y_i)=\check{y}_i$, $\varphi(p_j)=\check{p}_j$ for any $1\leq i\leq m$, $1\leq j\leq\nu$,*

(2) *$\check{u}\circ\varphi=u$.*

(3) *near every periodic orbit x, $\tilde{u}$ and $\check{\tilde{u}}\circ\varphi$ may differ by a canonical coordinate transformation* (26).

Definition 4.5. *Put*

$$Aut(u;((\Sigma,\mathbf{j}),\mathbf{y},\mathbf{p}))=\{\phi:\Sigma\to\Sigma|\phi \text{ is an automorphism satisfying (1), (2) and (3) in Definition 4.4}\}.$$

We call it the automorphism group of $(u;((\Sigma,\mathbf{j}),\mathbf{y},\mathbf{p}))$.

The following Lemma is obvious.

Lemma 4.6. *A J-holomorphic map $(u;((\Sigma,\mathbf{j}),\mathbf{y},\mathbf{p}))$ is stable if and only if $Aut(u;((\Sigma,\mathbf{j}),\mathbf{y},\mathbf{p}))$ is a finite group.*

Denote by $\mathcal{M}_A(M^+,C,g,m+\nu,\mathbf{y},\mathbf{p},(\mathbf{k},\mathfrak{e}))$ the moduli space of equivalence classes of all J-holomorphic curves in M^+ representing the homology class A and satisfying $(\mathbf{k},\mathfrak{e})$ boundary condition.

Fix $A\in H^2(M,L;\mathbb{Z})$, denote by $\mathcal{M}_A(M,L;g,m+\nu,\mathbf{y},\mathbf{p},\overrightarrow{\mu})$ the moduli space of equivalence classes of all J-holomorphic curves in M^+ representing the homology class A and satisfying $(\overrightarrow{\mu})$ boundary condition.

Lemma 4.7. *There is a constant $C>0$ depending on A and $\mathbf{k}$ such that for any $b=(u;(\Sigma,j),\mathbf{y},\mathbf{p})\in\mathcal{M}_A(M^+,C,g,m+\nu,\mathbf{y},\mathbf{p},(\mathbf{k},\mathfrak{e}))$ we have, over the cylinder end,*

$$E_\phi(u)+\int_\Sigma u^*d\lambda\leq C. \tag{47}$$

Proof. By the Stokes formula we have $\int_\Sigma u^*d\lambda \leq \sum_{i=1}^{\nu} k_i \cdot T_{\mathfrak{e}_i}$ over the cylinder end. Then the lemma follows from (46) $\Box$

Similarly, we also have

Lemma 4.8. *There is a constant $C>0$ depending on A and $\overrightarrow{\mu}$ such that for any $b=(u;(\Sigma,j),\mathbf{y},\mathbf{p}) \in \mathcal{M}_A(M,L;g,m+\nu,\mathbf{y},\mathbf{p},\overrightarrow{\mu})$ we have, over cylinder end,*

$$E_\phi(u) + \int_\Sigma u^*d\lambda \leq C. \tag{48}$$

We call $\mathcal{M}_A(M^+,C,g,m+\nu,\mathbf{y},\mathbf{p},(\mathbf{k},\mathfrak{e}))$ a holomorphic block in M^+.

Let $b=(u;(\Sigma,j),\mathbf{y},\mathbf{p}) \in \mathcal{M}_A(M^+,C,g,m+\nu,\mathbf{y},\mathbf{p},(\mathbf{k},\mathfrak{e}))$, $D_u : \mathcal{W}^{1,p,\alpha} \to L^{p,\alpha}$ be a Fredholm operator with $ind = dim(kerD_u) - dim(cokerD_u)$. Put

$$Ind^C = ind + 6(g-6) + 2(m+\nu).$$

The virtual dimension of $\mathcal{M}_A(M^+,C,g,m+\nu,\mathbf{y},\mathbf{p},(\mathbf{k},\mathfrak{e}))$ is Ind^C.

Similarly, let $b=(u;(\Sigma,j),\mathbf{y},\mathbf{p}) \in \mathcal{M}_A(M,L;g,m+\nu,\mathbf{y},\mathbf{p},\overrightarrow{\mu})$. Suppose that $\lim\limits_{s_j\to\infty} u(s_j,S^1) \subset \mathcal{F}_{i_j}$ and

$$[\pi(\lim_{s_j\to\infty} u(s_j,S^1))] = \mu_j, \;\; \forall\, 1 \leq j \leq \nu. \tag{49}$$

Then $D_u : \mathcal{W}^{1,p,\alpha} \to L^{p,\alpha}$ be a Fredholm operator with $ind = dim(kerD_u) - dim(cokerD_u)$. Put

$$Ind^L = ind + 6(g-6) + 2(m+\nu).$$

The virtual dimension of $\mathcal{M}_A(M,L;g,m+\nu,\mathbf{y},\mathbf{p},\overrightarrow{\mu})$ is Ind^L.

It is possible that there are finite many combinations such that (49) holds. In this case $\mathcal{M}_A(M,L;g,m+\nu,\mathbf{y},\mathbf{p},\overrightarrow{\mu})$ is the union of some $\mathcal{M}_A(M^+,C,g,m+\nu,\mathbf{y},\mathbf{p},.)$.

Remark 4.9. *As we mod the S^1 action on every periodic orbit of Reeb vector field, the situation is very similar to the symplectic cutting. For example, let $b=(u;\Sigma,j,p) \in \mathcal{M}_A(M^+,C;g,m+1,(k,\mathfrak{e}_1))$. Roughly speaking :*

We collapse the S^1-action on the orbit $x(kT_{\mathfrak{e}_1}t)$ at infinity to get a "manifold" $\widehat{M}^+$, locally , such that $Z_{\mathfrak{e}_1}$ is a "submanifold" of $\widehat{M}^+$, choose a local pseudo-Daubaux coordinate $(a,\vartheta,\mathbf{w})$ around $x(kT_{\mathfrak{e}_1}t)$. Our estimates (27),(28), (29) show that the puncture point p can be "removed", we get a J-holomorphic map $\bar{u}$ from Σ into $\widehat{M}^+$. The condition that u converges to a k-multiple periodic orbit at p is naturally interpreted as $\bar{u}$ being tangent to $Z_{\mathfrak{e}_1}$ at p with order k.

We will study the local geometry in other paper.

4.4 Holomorphic blocks in $\mathbb{R} \times \widetilde{M}$

Note that the space $\mathcal{M}_A(M^+,C;g,m+\nu,(\mathbf{k},\mathfrak{e}))$ is not large enough to compactify the moduli space of all J-holomorphic maps into M^+, we need consider $\mathcal{M}_A(\mathbb{R}\times\widetilde{M},C;(\mathbf{k}^-,\mathfrak{e}^-),(\mathbf{k}^+,\mathfrak{e}^+))$, which will be studied in this section.

Let $((\Sigma,\mathbf{j});\mathbf{y},\mathbf{p}^+,\mathbf{p}^-)$ be a connected semistable curve with m marked points $\mathbf{y}=(y_1,...,y_m)$ and $\nu^\pm$ puncture points $\mathbf{p}^+=(p_1^+,...,p_{\nu^+}^+)$, $\mathbf{p}^-=(p_1^-,...,p_{\nu^-}^-)$, and $u:\Sigma\to\mathbb{R}\times\widetilde{M}$ be a J-holomorphic map. Suppose that $u(z)$ converges to a $k_i^\pm \cdot T_{\mathfrak{e}_i^\pm}$-periodic orbit $x_{k_i^\pm} \subset \mathcal{F}_{\mathfrak{e}_i^\pm}$ as z tends to $p_i^\pm$.

Similar to the situation of relative invariants, there is a $\mathbb{R}$ action, which induces a $\mathbb{R}$-action on the moduli space of J-holomorphic maps. We need mod this action. Since there is a vector field X with $|X| = 1$ on $\widetilde{M}$, the Reeb vector field, there is a one parameter group φ_θ action on $\widetilde{M}$ generated by $-X$. In particular, there is a S^1-action on every periodic orbit, corresponding to the freedom of the choosing origin of S^1. Along every periodic orbit we have $L_X\lambda = 0$. Recall that by (3) of **Assumption** 2.4, $L_XJ = 0$ along every periodic orbit. We can mod this action.

Definition 4.10. *Two J-holomorphic maps $\Gamma = (u, (\Sigma, \mathbf{j}), \mathbf{y}, \mathbf{p}^+, \mathbf{p}^-)$ and $\check{\Gamma} = (\check{u}, (\check{\Sigma}, \check{\mathbf{j}}), \check{\mathbf{y}}, \check{\mathbf{p}}^+, \check{\mathbf{p}}^-)$ are called equivalent if there exists a diffeomorphism $\varphi : \Sigma \to \check{\Sigma}$ such that it can be lifted to bi-holomorphic isomorphisms $\varphi_{v\check{v}} : (\Sigma_v, j_v) \to (\check{\Sigma}_{\check{v}}, j_{\check{v}})$ for each component Σ_v of Σ, and*

- **(1)** *$\varphi(y_i) = \check{y}_i$, $\varphi(p_j^+) = \check{p}_j{}^+$, $\varphi(p_l^-) = \check{p}_l{}^-$ for any $1 \leq i \leq m$, $1 \leq j \leq \nu^+$, $1 \leq l \leq \nu^-$; u and $\check{u} \circ \varphi$ converges to the same periodic orbit $x_{k_i^\pm}$ at z tends to $p_i^\pm$;*
- **(2)** *$\check{a} \circ \varphi = a + C$, $\check{\tilde{u}} \circ \varphi = \tilde{u}$ for some constant C;*
- **(3)** *near every periodic orbit x, $\tilde{u}$ and $\check{\tilde{u}} \circ \varphi$ may differ by a canonical coordinate transformation* (26).

Definition 4.11. *A J-holomorphic map $(u; ((\Sigma, \mathbf{j}), \mathbf{y}, \mathbf{p}))$ is said to be stable if for each v one of the following conditions holds:*

- *(1). $\tilde{E}(u \circ \pi_{\Sigma_v}) \neq 0$,*
- *(2). Let val_v be the number of special points on Σ_v which are nodal points, marked points or puncture points. Then $val_v + 2g_v \geq 3$.*

Definition 4.12. *Put*

$$Aut(u; ((\Sigma, \mathbf{j}), \mathbf{y}, \check{\mathbf{p}}^+, \check{\mathbf{p}}^-)) = \{\phi : \Sigma \to \Sigma | \phi \text{ is an automorphism satisfying (1), (2) and (3) in Definition 4.10}\}.$$

We call it the automorphism group of $(u; ((\Sigma, \mathbf{j}), \mathbf{y}, \check{\mathbf{p}}^+, \check{\mathbf{p}}^-))$.

For any $A \in H^2(\mathbb{R} \times \widetilde{M}, \Re; \mathbb{Z})$ we define $d\lambda(A)$ as following: let $v : \mathbb{R} \times S^1 \to \mathbb{R} \times \widetilde{M}$ be a C^∞ map such that $[v(\mathbb{R} \times S^1)] = A$, we define $d\lambda(A) := \int_{\mathbb{R}\times S^1} v^*(d\lambda)$.

We fix $A \in H^2(\mathbb{R} \times \widetilde{M}, \Re; \mathbb{Z})$ and $\mathbf{k}^\pm = (k_1^\pm, ..., k_{\nu^\pm}^\pm)$, $\mathfrak{e}^\pm = (\mathfrak{e}_1^\pm, ..., \mathfrak{e}_{\nu^\pm}^\pm)$ satisfying

$$d\lambda(A) = \sum_{i=1}^{\nu^+} k_i^+ \cdot T_{\mathfrak{e}_i^+} - \sum_{i=1}^{\nu^-} k_i^- \cdot T_{\mathfrak{e}_i^-}. \tag{50}$$

We define $\mathcal{M}_A(\mathbb{R} \times \widetilde{M}, g, m + \nu^+ + \nu^-, (\mathbf{k}^-, \mathfrak{e}^-), (\mathbf{k}^+, \mathfrak{e}^+))$ to be the space of equivalence classes of all stable J-holomorphic maps in $\mathbb{R} \times \widetilde{M}$ representing A and converging to a $k_i^\pm \cdot T_{\mathfrak{e}_i^\pm}$-periodic orbit $x_{k_i^\pm} \subset \mathcal{F}_{\mathfrak{e}_i^\pm}$ as z tends to $p_i^\pm$. For any $(u, \Sigma, \mathbf{y}, \mathbf{p}^+, \mathbf{p}^-) \in \mathcal{M}_A(\mathbb{R} \times \widetilde{M}, g, m + \nu^+ + \nu^-, (\mathbf{k}^-, \mathfrak{e}^-), (\mathbf{k}^+, \mathfrak{e}^+))$, by Stoke's formula we have

$$d\lambda(A) = \int_\Sigma u^* d\lambda = \sum_{i=1}^{\nu^+} \lambda(k_i^+ x_i^+) - \sum_{i=1}^{\nu^-} \lambda(k_i^- x_i^-) = \sum_{i=1}^{\nu^+} k_i^+ \cdot T_{\mathfrak{e}_i^+} - \sum_{i=1}^{\nu^-} k_i^- \cdot T_{\mathfrak{e}_i^-}.$$

We call $\mathcal{M}_A(\mathbb{R} \times \widetilde{M}, C; g, m + \nu^+ + \nu^-, (\mathbf{k}^-, \mathfrak{e}^-), (\mathbf{k}^+, \mathfrak{e}^+))$ a holomorphic rubber block in $\mathbb{R} \times \widetilde{M}$.

Remark 4.13. *Let* $\Sigma_v \subset \Sigma$ *be a smooth connected component of* Σ*. Suppose that* $u(\Sigma_v)$ *lie in a compact set* $K \subset \mathbb{R} \times \widetilde{M}$*. Then every nodal point of* Σ_v *is a removable singular point. As* $\omega_\phi = d(\phi\lambda)$ *we conclude that* $E_\phi(u) \mid_{\Sigma_v} = 0$*, then* $u(\Sigma_v)$ *is one point, then* $\widetilde{E}(u) \mid_{\Sigma_v} = 0$*.*

4.5 Local coordinate system of holomorphic blocks

Let $b = (u, \Sigma, \mathbf{y}, \mathbf{p}^+, \mathbf{p}^-) \in \mathcal{M}_A(\mathbb{R}\times\widetilde{M}, C; g, m+\nu^+ +\nu^-, (\mathbf{k}^-, \mathfrak{e}^-), (\mathbf{k}^+, \mathfrak{e}^+))$, and $h \in \mathcal{W}^{1,p,\alpha}(\Sigma, u^*T(\mathbb{R}\times \widetilde{M}))$ such that $D_u(h) = 0$. Under the transformation

$$\hat{a} = a + C \tag{51}$$

and the canonical transformation of local pseudo-Daubaux coordinate system

$$\hat{\vartheta} = \vartheta + \vartheta_0 \tag{52}$$

h is invariant (see subsection §7.1.2, Case 2 below), so we can view h as a element in $T_b(\mathcal{M}_A(\mathbb{R} \times \widetilde{M}, C; m + \nu^+ + \nu^-, (\mathbf{k}^-, \mathfrak{e}^-), (\mathbf{k}^+, \mathfrak{e}^+))$. We assume that D_u is surjective. Let $e_1, ..., e_a$ be a base of $ker D_u$. Then for any $h \in ker D_u$ we have $h = \sum_i^a x_i e_i$. Let $w_1, ..., w_d$ be the local coordinates of $\mathcal{M}_{g,m+\nu^+ +\nu^-}$ around Σ, where $d = 6g - 6 + 2(m + \nu^+ + \nu^-)$. Then the $(w_1, ..., w_d, x_1, ..., x_a)$ is a local coordinate system of $\mathcal{M}_A(\mathbb{R} \times \widetilde{M}, C; m + \nu^+ + \nu^-, (\mathbf{k}^-, \mathfrak{e}^-), (\mathbf{k}^+, \mathfrak{e}^+))$.

Similarly, let $b = (u, \Sigma, \mathbf{y}, \mathbf{p}^+) \in \mathcal{M}_A(M^+, C; g, m + \nu, (\mathbf{k}, \mathfrak{e}))$, $h \in \mathcal{W}^{1,p,\alpha}(\Sigma, u^*T(M^+))$ with $D_u(h) = 0$, we view h as a element in $T_b(\mathcal{M}_A(M^+, C; g, m+\nu, (\mathbf{k}, \mathfrak{e}))$. Assume that D_u is surjective. We can choose a local coordinate system in $\mathcal{M}_{g,m+\nu}$ around Σ together with a coordinate system in $ker D_u$ as a local coordinate system of $\mathcal{M}_A(M^+, C; g, m + \nu, (\mathbf{k}, \mathfrak{e}))$.

5 Compactness theorems

We will use Li-Ruan's Compactification to our setting. Roughly speaking, the idea of Li-Ruan's Compactification is that firstly let the Riemann surfaces degenerate in Delingne-Mumford space and then let M^+ degenerate compatibly. We will give a detail description of the Li-Ruan's idea and give the definition of the stable compactification using language of graphs. This is one of the key parts of this paper.

5.1 Bubble phenomenon

The proofs of the following two theorems are standard (see [6]).

Theorem 5.1. *There is a constant* $\hbar > 0$ *such that for every* J*-holomorphic map* $u = (a, \widetilde{u}) : \mathbb{C} \to \mathbb{R} \times \widetilde{M}$ *with finite energy,*

(1) *if* $E_\phi(u) \neq 0$*, we have* $E_\phi(u) \geq \hbar$*;*

(2) *if* $\widetilde{E}(u) \neq 0$*, we have* $\widetilde{E}(u) \geq \hbar$*.*

Theorem 5.2. *There is a constant* $\hbar > 0$ *such that for every* J*-holomorphic map* $u \in \mathcal{M}_A(\mathbb{R} \times \widetilde{M}, C; g, m + \nu^\pm, (\mathbf{k}^-, \mathfrak{e}^-), (\mathbf{k}^+, \mathfrak{e}^+))$ *with finite energy, if* $\widetilde{E}(u) \neq 0$*, we have*

$$\widetilde{E}(u) = \sum_{i=1}^{\nu^+} k_i^+ - \sum_{i=1}^{\nu^-} k_i^- \geq \hbar.$$

Let $\Gamma^{(i)} = (u^{(i)}, \Sigma^{(i)}; \mathbf{y}^{(i)}, \mathbf{p}^{(i)}) \in \mathcal{M}_A(M^+, C, g, m+\nu, A, (\mathbf{k}, \mathfrak{e}))$ be a sequence. Then there is a constant $C > 0$ such that

$$\widetilde{E}(u^{(i)}) + E_\phi(u^{(i)}) < C \quad \forall\, i.$$

Suppose that $(\Sigma^{(i)}; \mathbf{y}^{(i)}, \mathbf{p}^{(i)})$ is stable and converges to $(\Sigma; \mathbf{y}, \mathbf{p})$ in $\overline{\mathcal{M}}_{g,m+\nu}$. On a neighborhood of $(\Sigma; \mathbf{y}, \mathbf{p})$ in $\overline{\mathcal{M}}_{g,m+\nu}$ we construct a smooth family of metrics on each $(\Sigma^{(i)}; \mathbf{y}^{(i)}, \mathbf{p}^{(i)})$ such that near each marked point the metric is the Euclidean metric on disc in $\mathbb{C}$, and near each nodal point the metric is the standard cylinder metric.

5.1.1 Bound of the number of singular points

Following McDuff and Salamon [23] we introduce the notion of singular points for a sequence $u^{(i)}$ and the notion of mass of singular points. We show that there is a constant $\hbar > 0$ such that the mass of every singular point is large than $\hbar$. Let q be a singular point and $q^{(i)} \in \Sigma^{(i)}$, $q^{(i)} \to q$. In case $u^{(i)}(q^{(i)}) \in M_0^+$ the argument is standard (see [23]). We only consider $\widetilde{E}$ over the cylinder end. Since the metrics near nodal points are the standard cylinder metrics, we have, in the cylinder coordinates $D_1(q^{(i)}) \subset \Sigma^{(i)} - \{nodal\ points\}$, where $D_1(q^{(i)}) = \{(s^{(i)}, t^{(i)}) \mid (s^{(i)} - s^{(i)}(q^{(i)}))^2 + (t^{(i)} - t^{(i)}(q^{(i)}))^2 \leq 1\}$. We identify $q^{(i)}$ with 0 and consider J-holomorphic maps $u^{(i)} : D_1(0) \to N$.

The proof of the following lemma is similar to Theorem 4.6.1 in [23]. We give the proof here for the reader's convenience.

Lemma 5.3. *Let $u^{(i)} : D_1(0) \to \mathbb{R} \times \widetilde{M}$ be a sequence of J-holomorphic maps with finite energy such that*

$$\sup_i \widetilde{E}(u^{(i)}) < \infty, \qquad |du^{(i)}(0)| \longrightarrow \infty, \ \ as \ \ i \to \infty.$$

Then there is a constant $\hbar > 0$ independent of $u^{(i)}$ such that, for every $\epsilon > 0$

$$\liminf_{i\to\infty} \widetilde{E}(u^{(i)}; D_\epsilon(0)) \geq \hbar. \tag{53}$$

Proof: Consider the function

$$F^{(i)}(z) = |du^{(i)}|(z) d^2(z, \partial D_1(0))$$

where $d(z, \partial D_1(0))$ denotes the distance from z to $\partial D_1(0)$ with respect to the standard Eucildean metrics. Obviously, $F^{(i)}$ attains its maximum at some interior point $q_i^\star \in D_1(0)$ and $\lim_{i\to\infty} F^{(i)}(q_i^\star) = \infty$. Set $\delta_i = \frac{1}{2} d(q_i^\star, \partial D_1(0))$. Then for any $q \in D_{\delta_i}(q_i^\star)$,

$$|du^{(i)}|(q) \leq 4|du^{(i)}|(q_i^\star) := 4A_i.$$

Consider the re-scaling sequence

$$v^{(i)}(z) = u^{(i)}\left(q_i^\star + \frac{z}{A_i}\right)$$

As $\lim_{i\to\infty} F_i(q_i^\star) = \infty$, we have $\lim_{i\to\infty} \delta_i A_i = \infty$. Then in $D_{\delta_i A_i}(0)$

$$\sup |dv^{(i)}| \leq 4, \quad |dv^{(i)}|(0) = 1, \quad \widetilde{E}(v^{(i)}, D_{\delta_i A_i}(0)) = \widetilde{E}(u^{(i)}, D_{\delta_i}(q_i^\star)).$$

By choosing a subsequence we conclude that v_i locally uniformly converges to a nonconstant J-holomorphic map with finite energy $v : \mathbb{C} \longrightarrow \mathbb{R} \times \widetilde{M}$. Then the lemma follows from Lemma 5.1. $\square$

Recall that N denotes one of M^+ and $\mathbb{R}\times\tilde{M}$. By Lemma 5.3 we conclude that the rigid singular points are isolated and the limit

$$m_\epsilon(q) = \lim_{i\to\infty} \widetilde{E}(u^{(i)}; D_{q^{(i)}}(\epsilon, h^{(i)}))$$

exists for every sufficiently small $\epsilon > 0$. The mass of the singular point q is defined to be

$$m(q) = \lim_{\epsilon\to 0} m_\epsilon(q).$$

Denote by $P \subset \Sigma$ the set of singular points for $u^{(i)}$, the double points and the puncture points. By Lemma 5.3 and (46), P is a finite set. By definition, $|du^{(i)}|_{h^{(i)}}$ is uniformly bounded on every compact subset of $\Sigma - P$. By a possible translation along $\mathbb{R}$ and passing to a subsequence we may assume that $u^{(i)}$ converges uniformly with all derivatives on every compact subset of $\Sigma - P$ to a J-holomorphic map $u : \Sigma - P \to N$. Obviously, u is a finite energy J-holomorphic map.

We need to study the behaviour of the sequence $u^{(i)}$ near each singular point for $u^{(i)}$. Let $q \in \Sigma$ be a rigid singular point for $u^{(i)}$. We have two cases.

(a) $q \in \Sigma - \{\mathit{nodal\ points}\}$. We consider J-holomorphic maps $u^{(i)} : D_1(0) \to N$.

(a-1) there are $\epsilon > 0$ and a compact set $K \subset N$ such that $u^{(i)}(D_\epsilon(q)) \subset K$.

(a-2) q is a nonremovable singularity.

(b) $q \in \{\mathit{nodal\ points}\}$. In this case a neighborhood of q is two discs $D_1(0)$ joint at 0, where $D_1(0) = \{|z|^2 \leq 1\}$.

For (a-1) we construct bubbles as usual for a compact symplectic manifold (see [23,25,28]). We call this type of bubbles (resp. bubble tree) the normal bubbles (resp. normal bubble tree).

5.1.2 Construction of the bubble tree for (a-2)

We use cylindrical coordinates $z = e^{-s-2\pi\sqrt{-1}t}$ and write

$$u^{(i)}(s,t) = (a^{(i)}(s,t), \widetilde{u}^{(i)}(s,t))$$

$$u(s,t) = (a(s,t), \widetilde{u}(s,t)).$$

Note that the graduate $|du^{(i)}|$ depends not only on the metric $<,>$ on N but also depends on the metric on $\Sigma^{(i)}$. The energy don't depend on the metric on $\Sigma^{(i)}$. To construct bubbling in present case it is more convenient to take the Euclidean metric $|dz|^2$ on the disk $D_1(0)$ and pullback to the coordinate (s,t) through $z = e^{-s-2\pi\sqrt{-1}t}$. Obviously, if q is a singular point with respect to the cylinder metric, then it is also a singular point with respect to the disk metric.

By Theorem 2.7, we have

$$\lim_{s\to\infty}\widetilde{u}(s,t)=x(kTt)$$

in $C^\infty(S^1)$, where $x(\ ,\ )$ is a kT-periodic orbit on $\widetilde{M}$. Choosing ϵ small enough we have

$$|m_\epsilon(q)-m(q)|\leq\frac{1}{10}\min\{\hbar,T\}$$

For every i there exists $\delta_i>0$ such that

$$\widetilde{E}(u^{(i)};D_{\delta_i}(0))=m(q)-\frac{1}{2}\min\{\hbar,T\},\tag{54}$$

where T is the minimal positive periodic. By definition of the mass $m(q)$, the sequence δ_i converges to 0. Since $u^{(i)}$ converges uniformly with all derivatives to u on any compact set of $D_\epsilon(0)-\{0\}$, δ_i must converge to 0. Put

$$\hat{s}^{(i)}=s+log\delta_i,\quad \hat{t}^{(i)}=t\ \ for\ |s_i^{(i)}|>2R_0.\tag{55}$$

$$\hat{a}^{(i)}=a-kT(-\log\delta_i).\tag{56}$$

Define the J-holomorphic curve $v^{(i)}(\hat{s},t)$ by

$$v^{(i)}(\hat{s},t)=(\hat{a}^{(i)}(\hat{s},t),\widetilde{v}^{(i)}(\hat{s},t))=\left(a^{(i)}(-\log\delta_i+\hat{s},t)-kT(-\log\delta_i),\widetilde{u}^{(i)}(-\log\delta_i+\hat{s},t)\right).\tag{57}$$

Lemma 5.4. *Suppose that 0 is a nonremovable singular point of u. Define the J-holomorphic map $v^{(i)}$ as above. Then there exists a subsequence (still denoted by $v^{(i)}$) such that*

(1) *The set of singular points $\{Q_1,\cdots,Q_d\}$ for $v^{(i)}$ is finite and tame, and is contained in the disc $D_1(0)=\{z\mid|z|\leq1\}$;*

(2) *The subsequence $v^{(i)}$ converges with all derivatives uniformly on every compact subset of $\mathbb{C}\backslash\{Q_1,\cdots,Q_d\}$ to a nonconstant J-holomorphic map $v:\mathbb{C}\backslash\{Q_1,\cdots,Q_d\}\to\mathbb{R}\times\widetilde{M}$;*

(3) $\widetilde{E}(v)>\frac{1}{3}\min\{\hbar,T\}$;

(4) $\widetilde{E}(v)+\sum_1^d m(Q_i)=m(0)$.

(5) $\lim_{s\to\infty}\widetilde{u}(s,t)=\lim_{\hat{s}\to-\infty}\widetilde{v}(\hat{s},t)$.

Proof: The proofs of **(1)**, **(2)** and **(4)** are standard (see [23]), we omit them here. The proof of **(5)** will be given in our next paper (see [19]). We only prove **(3)**.

(3) Note that

$$\left|\int_{\widetilde{u}^{(i)}(-\log\delta_i,S^1)}\lambda-\int_{\widetilde{u}^{(i)}(-\log\epsilon,S^1)}\lambda\right|=\widetilde{E}(u;-\log\epsilon\leq s\leq-\log\delta_i)=\frac{1}{2}\min\{\hbar,T\}.\tag{58}$$

Since $\lim_{s\to\infty}\widetilde{u}(s,t)=x(kTt)$ we have

$$\left|\int_{\widetilde{u}^{(i)}(-\log\epsilon,S^1)}\lambda-kT\right|\leq\frac{1}{8}\min\{\hbar,T\}.$$

when ϵ small enough and i big enough. Then by the locally uniform convergence of $v^{(i)}$

$$\widetilde{E}(v) \geq \left| kT - \int_{\widetilde{v}(0,S^1)} \lambda \right| \geq \frac{1}{3} \min\{\hbar, T\}. \tag{59}$$

□

We can repeat this again to construct bubble tree.

We introduce a terminology. Let Σ_1 and Σ_2 join at p, and $u_i : \Sigma_i \to \mathbb{R} \times \widetilde{M}$ a map. Choose holomorphic cylindrical coordinates $z_1 = (s_1, t_1)$ on Σ_1 and $z_2 = (s_2, t_2)$ on Σ_2 near p respectively. Suppose that

$$\lim_{s_1 \to -\infty} \widetilde{u_1}(s_1, t_1) = x_1(k_1 T_1 t_1)$$
$$\lim_{s_2 \to +\infty} \widetilde{u_2}(s_2, t_2) = x_2(k_2 T_2 t_2).$$

We say u_1 and u_2 converge to a same periodic orbit, if $k_1 = k_2$, and in the pseudo-Daubaux coordinates $(a, \vartheta, \mathbf{w})$, (see Lemma 2.5) $\mathbf{w}(x_1) = \mathbf{w}(x_2)$ (in this case $T_1 = T_2$ holds naturally).

5.1.3 Construction of bubble tree for (b)

Let $\Sigma = \Sigma_1 \wedge \Sigma_2$, where Σ_1 and Σ_2 are smooth Riemann surfaces of genus g_1 and g_2 joining at q. Let z_1 , z_2 be the local complex coordinates of Σ_1 and Σ_2 with $z_1(q) = z_2(q) = 0$. Recall that a neighborhood of $\Sigma_1 \wedge \Sigma_2 \in \overline{\mathcal{M}}_{g,m+\nu}$ is given by

$$z_1 z_2 = w = e^{-R - 2\pi\sqrt{-1}\tau}, \;\; w \in \mathbb{C},$$

where $R = 2lr$, $l \in \mathbb{Z}^+$. We will use (r, τ) as the local coordinates in the neighborhood of $\Sigma \in \overline{\mathcal{M}}_{g,m+\nu}$. Let

$$z_1 = e^{-s_1 - 2\pi\sqrt{-1}t_1}, \quad z_2 = e^{s_2 + 2\pi\sqrt{-1}t_2}.$$

(s_i, t_i) are called the holomorphic cylindrical coordinates near p. In terms of the holomorphic cylindrical coordinates we write

$$\Sigma_1 - \{p\} = \Sigma_{10} \bigcup \{[0, \infty) \times S^1\},$$

$$\Sigma_2 - \{p\} = \Sigma_{20} \bigcup \{(-\infty, 0] \times S^1\}.$$

$$s_1 = s_2 + 2lr \tag{60}$$
$$t_1 = t_2 + \tau. \tag{61}$$

Then $\Sigma_{(r)} \to \Sigma_1 \wedge \Sigma_2$ as $r \to \infty$ in the $\overline{\mathcal{M}}_{g,m+\nu}$.

We consider the case that $u^{(i)} : \Sigma^{(i)} \to N$ is a sequence of J-holomorphic maps, where $\Sigma^{(i)} := \Sigma_{(r^{(i)})}$, $q^{(i)} \in \Sigma^{(i)}$, $q^{(i)} \to q$.

Without loss of generality we assume that there is a subsequence, still denoted by i, such that $|s_1^{(i)}(q^{(i)})| \leq lr^{(i)}$. In this case we may identify a neighborhood of $q^{(i)}$ in $\Sigma^{(i)}$ with $D_1(0) \setminus D_{\epsilon^{(i)}}(0) \subset \Sigma_1$, where $\lim_i \epsilon^{(i)} = 0$. The sequence $u^{(i)}$ is considered to be a sequence of J-holomorphic maps from

$D_1(0) \setminus D_{\epsilon^{(i)}}(0)$ into N, and $q^{(i)} \in D_1(0) \setminus D_{\epsilon^{(i)}}(0)$, $q^{(i)} \to 0$. In terms of the cylinder coordinates we can identify the coordinates $(s^{(i)}, t^{(i)})$ in $\Sigma^{(i)}$ with the coordinates (s_1, t_1) of Σ_1.

By Theorem 2.7, we have

$$\lim_{s_1 \to \infty} \widetilde{u}(s_1, t_1) = x(kTt_1)$$

in $C^\infty(S^1)$, where $x(\ ,\ )$ is a kT-periodic orbit on $\widetilde{M}$. Choosing ϵ small enough we have

$$|m_\epsilon(q) - m(q)| \leq \frac{1}{10}\min\{\hbar, T\}$$

For every i there exists $\delta_i > 0$ such that

$$\widetilde{E}(u^{(i)}; D_{\delta_i}(0)) = m(q) - \frac{1}{2}\min\{\hbar, T\}, \tag{62}$$

where T is the minimal positive periodic. We choose a smooth conformal transformation $\psi^{(i)} : \Sigma^{(i)} \to \Sigma^{(i)}$ such that

$$\psi^{(i)}|_{\mathfrak{D}(R_0)} = I, \tag{63}$$

$$z^{(i)} = \delta_i \hat{z}^{(i)} = \delta_i e^{-\hat{s}^{(i)} - 2\pi\sqrt{-1}\hat{t}^{(i)}}, \quad for\ |s^{(i)}| > 2R_0 \tag{64}$$

where $(\hat{s}^{(i)}, \hat{t}^{(i)}) = \psi_i(s^{(i)}, t^{(i)})$, i.e.,

$$\hat{s}^{(i)} = s^{(i)} + log\delta_i = s_1 + log\delta_i, \qquad \hat{t}^{(i)} = t^{(i)} = t_1 \ \ for\ |s_i^{(i)}| > 2R_0. \tag{65}$$

We call coordinates rescalling of Riemann surface a $\mathcal{D}$-rescaling. In our present case we need not only a $\mathcal{D}$-rescaling, but also a translation along $\mathbb{R}$, called a $\mathcal{T}$-rescaling. We call such composition a $\mathcal{DT}$-rescaling. Put

$$\hat{a}^{(i)} = a - kT(-\log\delta_i). \tag{66}$$

Define the J-holomorphic curve $v^{(i)}(\hat{s}, t)$ by

$$v^{(i)}(\hat{s}, t) = (\hat{a}^{(i)}(\hat{s}, t), \widetilde{v}^{(i)}(\hat{s}, t)) = \tag{67}$$

$$\left(a^{(i)}(-\log\delta_i + \hat{s}, t) - kT(-\log\delta_i), \widetilde{u}^{(i)}(-\log\delta_i + \hat{s}, t)\right).$$

By the same argument as in subsection §5.1.2 we construct a bubble S^2 with $E(v)\mid_{S^2} > \frac{1}{3}\min\{\hbar, T\}$ or $\widetilde{E}(v)\mid_{S^2} > \frac{1}{3}\min\{\hbar, T\}$, inserted between Σ_1 and Σ_2. The same results as Lemma 5.4 still hold. We can repeat this again to construct bubble tree.

5.1.4 For the case of genus 0

Let $\Gamma^{(i)} = (u^{(i)}, \Sigma^{(i)}; \mathbf{y}^{(i)}, \mathbf{p}^{(i)}) \in \mathcal{M}_A(M^+, C, g, m+\nu, \mathbf{y}, \mathbf{p}, (\mathbf{k}, \mathfrak{c}))$ be a sequence. Let $\Sigma^{(i)} = \bigcup_{v=1}^N \Sigma_v^{(i)}$. Assume that there is one component of $\Sigma^{(i)}$ that has genus 0 and is unstable. Let $\Sigma_1^{(i)}$ is such a component. We identify $\Sigma_1^{(i)}$ with a sphere S^2, and consider $u^{(i)} : S^2 \to N$. We discuss several cases:

1). $u^{(i)}\mid_{S^2}$ has no singular point. Then $\|\nabla u^{(i)}\mid_{S^2}\|$ are uniformly bounded above. As $(u^{(i)}, S^2; \mathbf{y}^{(i)}, \mathbf{p}^{(i)})$ is stable, $E(u^{(i)})\mid_{S^2} \geq \hbar$ or $\widetilde{E}(u^{(i)})\mid_{S^2} \geq \hbar$. Then $u^{(i)}\mid_{S^2}$ locally uniformly converges to $u\mid_{S^2}$ with $E(u)\mid_{S^2} \geq \hbar$ or $\widetilde{E}(u)\mid_{S^2} \geq \hbar$, so $(u; S^2)$ is stable.

2). S^2 joints with $\Sigma_2^{(i)}$ at a nodal point q. Then the number of the special points, including the marked points and the puncture points, is ≤ 1.

2a) There is one singular point p and one special point $\mathcal{X}$.

(2a-1) $p \neq q$ and $p \neq \mathcal{X}$,

(2a-2) $p = \mathcal{X}$ or $p = q$

2b) There are two singular points p_1, p_2.

(2b-1) $p_1 \neq q$ and $p_2 \neq q$,

(2b-2) $p_1 = q$ or $p_2 = q$.

2c) There is only one singular point p and no special point.

3). S^2 joint with $\Sigma_2^{(i)}$ and $\Sigma_3^{(i)}$ at q_1 and q_2 respectively. There is only one singular point p.

(3a) $p \neq q_1$ and $p \neq q_2$,

(3b) $p = q_1$ or $p = q_2$,

For the cases **(2a-1)**, **(2b-1)**, **3a)** we construct bubble tree at singular points as in subsection §5.1.2 to get a stable map (u, S^2). For the cases **(2a-2)**, **(2b-2)**, **2c)** and **(3b)** we forget the map $u^{(i)} |_{S^2}$ and contract S^2 as a point, then we construct a bubble as in subsection subsection §5.1.3. By (**3**) of Lemma 5.4 we get a stable map (v, S^2). We can repeat the procedure to construct bubble tree.

5.2 $\mathcal{T}$- rescaling

Let $\Gamma^{(i)} = (u^{(i)}, \Sigma^{(i)}; \mathbf{y}^{(i)}, \mathbf{p}^{(i)}) \in \mathcal{M}_A(M^+, C, g, m+\nu, \mathbf{y}, \mathbf{p}, (\mathbf{k}, \mathfrak{c}))$ be a sequence. If there is some Σ_v of genus 0, we treat it as in subsection 5.1.4. In the following we assume that $(\Sigma^{(i)}; \mathbf{y}^{(i)}, \mathbf{p}^{(i)})$ is stable and converges to $(\Sigma; \mathbf{y}, \mathbf{p})$ in $\overline{\mathcal{M}}_{g,m+\nu}$.

Note that we have fixed the degeneration of Riemann surfaces now, we are concern with the $\mathcal{T}$-rescaling. We explain our procedure of $\mathcal{T}$-rescaling. We discuss maps into M^+, for maps into $\mathbb{R} \times \widetilde{M}$ the situation are the same.

Let Σ be a Riemann surface with ν puncture points $\mathbf{p} = (p_1, ..., p_\nu)$. We choose holomorphic coordinates (s_i, t_i) near p_i. Let $u : \overset{\circ}{\Sigma} \to M^+$ be a J-holomorphic map. Denote $u = (a, \tilde{u})$. Suppose that

$$\lim_{|s_j| \to \infty} \widetilde{u}(s_j, t_j) = x(k_j T_{\mathfrak{c}_j} t_j)$$

in $C^\infty(S^1)$ for some $k_j T_{\mathfrak{c}_j}$-periodic orbit $x(k_j T_{\mathfrak{c}_j} t)$. For each j we choose a local pseudo-Darboux coordinates $(a_j, \vartheta_j, \mathbf{w_j})$ near p_j such that $x(k_j T_{\mathfrak{c}_j} t_j) = (k_j T_{\mathfrak{c}_j} t_j + \vartheta_{j0}, 0)$ for some constant ϑ_j. Note that for any $0 < l, j \leq \nu$,

$$a_l = a_j + C_{lj},$$

where C_{lj} are constants.

For simplicity, we consider the case as in Figure 1, the other cases is similar. Let $\Sigma = \bigcup_{j=1}^{6} \Sigma_j$, where each Σ_j is a smooth connected component. Let $T_j \subset \Sigma_j - P$ be compact sets, where $P \subset \Sigma$

denotes the set of singular points for $u^{(i)}$, the double points and the puncture points. Denote $I = \{1, \cdots, 6\}$. We write

$$u^{(i)}(z) = (a^{(i)}(z), \widetilde{u}^{(i)}(z)).$$



Figure 1:

For each $j \in I$, $T_j^{(i)} \subset \Sigma^{(i)}$ converges to T_j. We fix points $z_j \in T_j$. Suppose that $z_j^{(i)} \in T_j^{(i)}$ is a sequence of points which converges to z_j. Without loss of generality, we assume that

(1) $\left|a^{(i)}(z_1^{(i)})\right| \leq \min_{j \in I \setminus \{1\}} \left|a^{(i)}(z_j^{(i)})\right|, \ \forall i,$

(2) $\sup\limits_i \left|a^{(i)}(z_1^{(i)})\right| < \infty,$

Restricting to T_1, $u^{(i)}$ uniformly converges to a map u_1. For any compact $K \subset \Sigma_1 - P$, $u^{(i)}$ also converges to u_1. Then u is naturally defined on $\Sigma_1 - P$.

For Σ_2 we consider two different cases:

Case 1. $\sup_i \left|a^{(i)}(z_2^{(i)})\right| < \infty$. Restricting to T_2, $u^{(i)}$ uniformly converges to a map u_2. Then u_2 is naturally defined on $\Sigma_2 - P$. Then $(u_1; \Sigma_1, \mathbf{y}_1, \mathbf{p}_1)$ and $(u_2; \Sigma_2, \mathbf{y}_2, \mathbf{p}_2)$ are belong to the same holomorphic block in M^+.

Case 2. $\lim\limits_{k \to \infty} a^{(i)}(z_2^{(i)}) = \infty$. In this case we take a coordinate transformation:

$$(a^*)^{(i)} = a - a^{(i)}(z_2^{(i)}). \tag{68}$$

Define

$$(u^*)^{(i)}(z) = ((a^*)^{(i)}(z), (\tilde{u}^*)^{(i)}(z)), \qquad (\tilde{u}^*)^{(i)}(z) = \tilde{u}^{(i)}(z). \tag{69}$$

Note that $|du^{(i)}|$ is invariant under translation along $\mathbb{R}$. As above, restricting to $T_2^{(i)}$, $(u^*)^{(i)}$ locally uniformly converges to a map $u^* = (a^*, \tilde{u}^*) : T_2 \longrightarrow \mathbb{R} \times \widetilde{M}$ with $a^*(z_2) = 0$, which extends to $u^* : \Sigma_2 - P \longrightarrow \mathbb{R} \times \widetilde{M}$. If $\lim\limits_{\epsilon \to 0} m_\epsilon(q_2) \geq \hbar$, then we can construct bubbles as in subsection §5.1.3. Otherwise, u and u^* converges to the same periodic orbit as $z \to q_2$. Note that, near q_2, in terms of the local cylinder coordinates the degeneration is given by

$$(s^*)^{(i)} = s - 2lr^{(i)}, \qquad (t^*)^{(i)} = t - \tau^{(i)}. \tag{70}$$

In local pseudo-Dauboux coordinate system $(a, \vartheta, \mathbf{w})$ and the cylinder coordinates of Riemann surface near q_2 we write

$$u(s,t) = (a(s,t), \widetilde{u}(s,t)), \quad u^*(s^*,t^*) = (a^*(s^*,t^*), \widetilde{u}^*(s^*,t^*)). \tag{71}$$

Then

$$\lim_{|s|\to\infty} \widetilde{u}(s,t) = x(kTt), \qquad \lim_{|s^*|\to\infty} \widetilde{u}^*(s^*,t^*) = x(kTt^* + \vartheta_0^*) \tag{72}$$

in $C^\infty(S^1)$ for some kT-periodic orbit $x(kTt)$.

We call $u^* : \Sigma_2 \longrightarrow \mathbb{R} \times \widetilde{M}$ a rubber component.

By the same argument we can construct maps on $\Sigma_j - P, j \geq 3$.

Remark 5.5. *We use notations above. Suppose that* $\sup_i \left| a_1^{(i)}(z_1^{(i)}) \right| \to \infty$ *and there is only one singular point* $p \in \Sigma_1$ *with*

$$p^{(i)} \to p, \ \sup_i \left| u^{(i)}(p^{(i)}) \right| < \infty.$$

We re-scale as above and construct bubble tree as usual. Then all main part of $u(\Sigma)$ *are rubber component and there is a bubble tree with* $u(p) \in M^+$*.*

Remark 5.6. *For Figure 1, it is possible that some* $u^{(i)}(T_j^{(i)}) \subset M^+$ *for some* $j \geq 2$*. We assume that*

$$u^{(i)}(T_1^{(i)}) \subset M^+, \quad u^{(i)}(T_6^{(i)}) \subset M^+,$$

and

$$\sup_{i\to\infty} |a^{(i)}(z_1^{(i)})| < \infty, \quad \sup_{i\to\infty} |a^{(i)}(z_6^{(i)})| < \infty. \tag{73}$$

In this case there is a relation between $a^{(i)}(z_j^{(i)})$, $1 \leq j \leq 6$*. To see this and to simplify notations, we omit the index* (i) *and let*

$$l_1 = a(z_3) - a(z_1), \quad l_2 = a(z_4) - a(z_3),$$
$$l_3 = a(z_5) - a(z_4), \quad l_4 = a(z_6) - a(z_5).$$

Then

$$l_1 + l_2 + l_3 + l_4 = a(z_6) - a(z_1). \tag{74}$$

This means that $l_1 + l_2 + l_3 + l_4$ *is a* $\mathcal{T}$ *re-scaling invariant. By* (73)*, there is only three independent parameter.*

If we start from Σ_6 *to re-scale by the above procedure, we get the same result. Similarly, we start from any component* Σ_i*,* $i \geq 3$*, we get also the same result. This suggests a alternative way to do* $\mathcal{T}$*-re-scaling for* Σ*: we first re-scale for both* Σ_1 *and* Σ_6 *by*

$$\breve{a}^{(i)}(z) = a^{(i)}(z) - a^{(i)}(z_3) = a^{(i)}(z) - l_1^{(i)} - a^{(i)}(z_1),$$
$$\acute{a}^{(i)}(z) = a^{(i)}(z) - a^{(i)}(z_5) = a^{(i)}(z) + l_4^{(i)} - a^{(i)}(z_6),$$

then we re-scale for both Σ_3 *and* Σ_5 *by*

$$a^{\bullet^{(i)}}(z) = \breve{a}^{(i)}(z) - \breve{a}^{(i)}(z_4) = \breve{a}^{(i)}(z) - l_2^{(i)},$$
$$a^{\bullet^{(i)}(z)} = \acute{a}^{(i)}(z) - \acute{a}^{(i)}(z_4) = \acute{a}^{(i)}(z) + l_3^{(i)}.$$

In view of (75) *we find that the above re-scalings are equivalent.*

In the following we consider the singular points. Let $\tilde{q}$ be a singular point of $u^{(i)}$. We discuss two cases:

Case a. $\tilde{q}$ is not a node of Σ. In this case we construct bubble tree as usual.

Case b. $\tilde{q}$ is a node of Σ. Suppose that $\tilde{q} = q_1$. We construct bubble as in subsection §5.1.3 to get S^2, inserted between Σ_1 and Σ_3, with $\widetilde{E}(v)\mid_{S^2} > \frac{1}{3}\min\{\hbar, T\}$. Denote $q_1' = \Sigma_1 \cap S^2$, $q_1 = \Sigma_3 \cap S^2$.

We fix a point $\mathfrak{z}$ in the compact set of $S^2 - \{q_1, q_1'\}$. Let $\mathfrak{z}^{(i)} \in \Sigma^{(i)}$ such that $\mathfrak{z}^{(i)} \to \mathfrak{z}$. Then $\Sigma' = \Sigma \cup S^2$ is a semi-stable curve and there exists a fixed degeneration $\Sigma^{(i)} \to \Sigma'$.

Consider the coordinate transformations

$$\hat{a}^{(i)} = a - kT(-\log \delta_i),$$

and

$$(a^{\circ})^{(i)} = \hat{a}^{(i)} - \hat{a}^{(i)}(z_3^{(i)}).$$

Note that for any $z \in \Sigma_3$

$$\hat{a}^{(i)}(z) - \hat{a}^{(i)}(z_3^{(i)}) = a^{(i)}(z) - a^{(i)}(z_3^{(i)}).$$

Then we have

$$(a^{\circ})^{(i)}(z) = (\check{a})^{(i)}(z).$$

To simplify notations, we omit the index (i) and let

$$l_1' = a(\mathfrak{z}) - a(z_1), \quad l_1^{\star} = a(z_3) - a(\mathfrak{z}), \quad l_2 = a(z_4) - a(z_3),$$

$$l_3 = a(z_5) - a(z_4), \quad l_4 = a(z_6) - a(z_5).$$

Then

$$l_1' + l_1^{\star} + l_2 + l_3 + l_4 = a(z_6) - a(z_1). \tag{75}$$

We conclude that the $\mathcal{T}$-rescalling based on $\Sigma^{(i)} \to \Sigma$ and the $\mathcal{T}$-rescalling based on $\Sigma^{(i)} \to \Sigma'$ are equivalent.

5.3 Equivalent $\mathcal{DT}$- rescaling

We consider the example in subsection 5.2 (see Figure 1). For simplicity we only consider the degeneration of $\Sigma_{(r)}$ at q_2 and consider the **Case 2** here. Let $q^{(i)} \in \Sigma^{(i)}$, $q^{(i)} \to q_2$. Suppose that there exists a constant $N > 0$ such that

$$\lim_{i\to\infty} \sup \tilde{E}(u^{(i)}; N-1 \le s \le 2lr^{(i)} - N + 1) \le \frac{1}{2}\min\{\hbar, T\}. \tag{76}$$

We have two $\mathcal{DT}$-rescaling:

A. The degeneration of Riemann surfaces in the Delingne-Mumford space, together with the $\mathcal{T}$-re-scaling in Section §5.2, we have a $\mathcal{DT}$-rescaling given by (68), (69) and (70)

B. There is another re-scaling as following. Assume that $\Sigma^{(i)}$ degenerate at q_2 with the formula (70). We choose a new coordinate system $\check{a}^{(i)}$ by

$$\check{a}^{(i)} = a - 2kTlr^{(i)}. \tag{77}$$

Define

$$\breve{u}^{(i)}(z) = (\breve{a}^{(i)}(z), \breve{\tilde{u}}^{(i)}(z)) = (a^{(i)}(z) - 2kTlr^{(i)}, \tilde{u}^{(i)}(z)). \tag{78}$$

Then the two coordinate systems $\breve{a}^{(i)}$ and $(a^*)^{(i)}$ satisfies

$$\breve{a}^{(i)} = (a^*)^{(i)} + a^{(i)}(z_2^{(i)}) - 2kTlr^{(i)}. \tag{79}$$

Choose two sequences of points $\mathfrak{z}_1^{(i)}, \mathfrak{z}_2^{(i)}$ such that $(s,t)(\mathfrak{z}_1^{(i)}) = (N, 0)$ and $((s^*)^{(i)}, (t^*)^{(i)})(\mathfrak{z}_2^{(i)}) = (-N, 0)$. Obviously, $\mathfrak{z}_j^{(i)} \to \mathfrak{z}_j \in \Sigma_j - P$, $j = 1, 2$. It follows from the convergence of $u^{(i)}$ and $(u^*)^{(i)}$ on compact sets that

$$|a^{(i)}(\mathfrak{z}_1^{(i)}) - kTs(\mathfrak{z}_1^{(i)})| \le C_2, \quad |a^{(i)}(\mathfrak{z}_2^{(i)}) - a^{(i)}(z_2^{(i)})| = |(a^*)^{(i)}(\mathfrak{z}_2^{(i)}) - (a^*)^{(i)}(z_2^{(i)})| \le C_3. \tag{80}$$

By the same argument of (**5**) in Lemma 5.4 we have

$$|a^{(i)}(\mathfrak{z}_2^{(i)}) - kTs(\mathfrak{z}_2^{(i)})| \le C_3'.$$

By (70) we have

$$s(\mathfrak{z}_2^{(i)}) - 2lr^{(i)} = s^*(\mathfrak{z}_2^{(i)}) = -N.$$

Then

$$|a^{(i)}(\mathfrak{z}_2^{(i)}) - 2kTlr^{(i)}| \le |a^{(i)}(\mathfrak{z}_2^{(i)}) - kTs(\mathfrak{z}_2^{(i)})| + |kTs(\mathfrak{z}_2^{(i)}) - 2kTlr^{(i)}| \le C_3' + kTN \le C_4. \tag{81}$$

It follows from (79), (80) and (81) that for any point $z \in \Sigma_2 - P$ we have

$$|\breve{a}^{(i)}(z) - (a^*)^{(i)}(z)| \le C \tag{82}$$

for some positive constant C.

We call two $\mathcal{DT}$-rescalings are equivalent if there exists a constant $C > 0$ independent of i such that (82) holds. We have proved

Lemma 5.7. *Let $\Sigma = \Sigma_1 \wedge \Sigma_2$, where Σ_1 and Σ_2 are smooth Riemann surfaces of genus g_1 and g_2 joining at q. Assume that $\Sigma^{(i)}$ is a sequence of smooth Riemann surface which converges to Σ in the Delingne-Mumford space as $i \to \infty$. Suppose that*

$$\lim_{i\to\infty} \sup E(u^{(i)}; D_\epsilon(q)) \le \frac{1}{2}\min\{\hbar, T\}, \tag{83}$$

and restricting on $\Sigma_2 \setminus D_\epsilon(q)$ there is no singular points of $u^{(i)}$. Then the two $\mathcal{DT}$-rescalings **A** *and* **B** *are equivalent.*

The above discussion can be immediately generalized to the case of several nodal points.

Lemma 5.8. *Let $\Sigma = \Sigma_1 \wedge \Sigma_2$, where Σ_1 and Σ_2 are smooth Riemann surfaces of genus g_1 and g_2 joining at $q_1, \cdots, q_\nu$. Assume that $\Sigma^{(i)} = \Sigma_1 \#_{(\mathbf{r}^{(i)})} \Sigma_2$ be a sequence of smooth Riemann surface which converges to Σ in the Delingne-Mumford moduli space as $i \to \infty$. Let $u^{(i)} : \Sigma^{(i)} \to M^+$ be a sequence J-holomorphic maps. Suppose that for any $j \in \{1, \cdots, \nu\}$,*

$$\lim_{i\to\infty} \sup E(u^{(i)}; D_\epsilon(q_j)) \le \frac{1}{2}\min\{\hbar, T\}, \tag{84}$$

and restricting on $\Sigma_2 - \bigcup_{j=1}^{\nu} D_\epsilon(q_j)$ there is no singular points of $u^{(i)}$. Then the $\mathcal{DT}$-rescaling of type **A** *and any $\mathcal{DT}$ re-scalings of type* **B** *are equivalent.*

5.4 Procedure of re-scaling

To sum up, for any sequence $\Gamma^{(i)} = (u^{(i)}, \Sigma^{(i)}; \mathbf{y}^{(i)}, \mathbf{p}^{(i)}) \in \mathcal{M}_A(M^+, C, g, m+\nu, (\mathbf{k}, \mathfrak{e}))$ our procedure is following:

(1) If there is some Σ_v of genus 0, we treat it as in subsection 5.1.4. In the following we assume that $(\Sigma^{(i)}; \mathbf{y}^{(i)}, \mathbf{p}^{(i)})$ is stable and converges to $(\Sigma; \mathbf{y}, \mathbf{p})$ in $\overline{\mathcal{M}}_{g,m+\nu}$.

(2) By Lemma 5.3 the number of singular points of Σ is finite. Denote by $P \subset \Sigma$ the set of singular points for $u^{(i)}$, the nodal points and the puncture points.

(3) We first find a component Σ_k of Σ, for example Σ_1 in Figure 1, such that

$$\left|a_1^{(i)}(z_1^{(i)})\right| \leq \min_{j \in I \setminus \{1\}} \left|a_j^{(i)}(z_j^{(i)})\right|, \ \ \forall i.$$

Without loss of generality we suppose that $\sup_i \left|a_1^{(i)}(z_1^{(i)})\right| < \infty$, that is, $u^{(i)}(z_1^{(i)}) \subset M^+$. Find a set $J \subset I$ such that $j \in J$ if and only if $u(\Sigma_j) \subset M^+$, for example Σ_1 and Σ_6 in Figure 1, i.e., $J = \{1, 6\}$. Let $\Sigma^{1\bigstar} = \Sigma - \bigcup_{j \in J} \Sigma_j$. $\Sigma^{1\bigstar}$ may have several connected components. For every connected component of $\Sigma^{1\bigstar}$ we do $\mathcal{T}$-rescaling independently as in §5.2.

We repeat the procedure. Finally we will stop after finite steps.

(4) Then we construct bubble tree for every singular point independently to get Σ', where Σ' is obtained by joining chains of $\mathbf{P^1}s$ at some double points of Σ to separate the two components, and then attaching some trees of $\mathbf{P^1}$'s. Then we have

(a) for every nodal point $q \in \mathfrak{N}$ there is a neighborhood $D_\epsilon(q)$ so that

$$\lim_{i \to \infty} \sup E(u^{(k)}; D_\epsilon(q)) \leq \frac{\hbar}{2},$$

where $\mathfrak{N}$ denotes the set of all nodal points of Σ';

(b) restricting on $\Sigma' \setminus \bigcup_{q \in \mathfrak{N}} D_\epsilon(q)$ there is no singular points of $u^{(i)}$. $\Box$

For every sequence $\Gamma^{(i)} = (u^{(i)}, \Sigma^{(i)}; \mathbf{y}^{(i)}, \mathbf{p}^{(i)}) \in \mathcal{M}_A(M^+, C, g, m+\nu, (\mathbf{k}, \mathfrak{e}))$, using our procedure we get $\Gamma = (u, \Sigma', \mathbf{y}, \mathbf{p})$, where

(A-1) Σ' is obtained by joining chains of $\mathbf{P^1}s$ at some double points of Σ to separate the two components, and then attaching some trees of $\mathbf{P^1}$'s. Σ' is a connected curve with normal crossings. We call components of Σ principal components and others bubble components.

(A-2) $u : \Sigma' \to (M^+)'$ is a continuous map, where $(M^+)'$ is obtained by attaching some $\mathbb{R} \times \widetilde{M}$ to M^+. Let Σ_1 be a connected component of Σ' and $u|_{\Sigma_1} : \Sigma_1 \to \mathbb{R} \times \widetilde{M}$. We call $(u; \Sigma_1)$, modulo the translations $\mathbb{R} \times \widetilde{M}$ and the S^1-action on periodic orbits, a rubber component.

(A-3) If we attach a tree of $\mathbf{P^1}$ at a marked point y_i or a puncture point p_i, then y_i or p_i will be replaced by a point different from the intersection points on a component of the tree. Otherwise, the marked points or puncture points do not change;

(A-4) Let val_v be the number of points on Σ_v which are nodal points or marked points or puncture points. In case $u(\Sigma_v) \subset M^+$, if $u|_{\Sigma_v}$ is constant then $val_v + 2g_v \geq 3$; in case $u : \Sigma_v \to \mathbb{R} \times \widetilde{M}$, if $\tilde{u}|_{\Sigma_v}$ is constant then $val_v + 2g_v \geq 3$;

(A-5) u converges exponentially to $(k_1 T_{\mathfrak{c}_1}, \cdots, k_\nu T_{\mathfrak{c}_\nu})$ periodic orbits $(x_{k_1}, ..., x_{k_\nu})$ as the variable tends to the puncture $(p_1, ..., p_\nu)$; more precisely, u satisfies (27)-(29);

(A-6) The restriction of u to each component of Σ' is J-holomorphic. Let q be a nodal point of Σ' . Suppose q is the intersection point of Σ_v and Σ_w. If q is a removable singular point of u, then u is continuous at q; If q is a nonremovable singular point of u, then $u|_{\Sigma_v}$ and $u|_{\Sigma_w}$ converge exponentially to the same periodic orbit on $\widetilde{M}$ as the variables tend to the nodal point q.

5.5 Weighted dual graph with a oriented decomposition

It is well-known that the moduli space of stable maps in a compact symplectic manifold has a stratification indexed by the combinatorial type of its decorated dual graph. In this section we generalizes this construction to our setting and in the next section we state Li-Ruan's compatification by using weighted dual graphs.

Let G be a graph. Denote $G = (V(G), E(G))$, where $V(G)$ is a finite nonempty set of vertices and $E(G)$ is a finite set of edges. Suppose that $V = \{v_1, ..., v_N\}$. Given a partition of $\{1, 2, ..., N\}$

$$\mathfrak{d} : \{1, \cdots, N\} = (\cup_{i=1}^{c} I_i) \bigcup (\cup_{\alpha=1}^{d} J_\alpha),$$

it induces a decomposition of V, still denoted by $\mathfrak{d}$,

$$\mathfrak{d} : V = \mathfrak{A} \bigcup \mathfrak{B},$$

where

$$\mathfrak{A} = \bigcup_{i=1}^{c} W_{I_i}, \quad \mathfrak{B} = \bigcup_{\alpha=1}^{d} W_{J_\alpha}, \quad W_{I_i} = \{v_k | k \in I_i\}, \quad W_{J_\alpha} = \{v_k | k \in J_\alpha\}.$$

Obviously, $\mathfrak{A} \bigcap \mathfrak{B} = \emptyset$. Every subset W_{I_i} (resp. W_{J_α}) determines a induced subgraph G_{I_i} (resp. G_{J_α}) of G.

Assumption 1. For any $W_{I_i}, W_{I_j} \subset \mathfrak{A}, i \neq j$, there is no edge in $E(G)$ connecting G_{I_i} and G_{I_j}.

Denote by $R(G)$ all the edges which connect two subgraphs above. Obviously,

$$R(G) = E(G) - (\cup_{i=1}^{c} E(G_{I_i})) \bigcup (\cup_{\alpha=1}^{d} E(G_{J_\alpha})).$$

Let $\ell \in R(G)$ be an edge connecting v_i and v_j. We give an orientation to ℓ, denoted by

$$\overrightarrow{\ell} : v_i \xrightarrow{\ell} v_j. \tag{85}$$

Sometimes we denote simply by $v_i \xrightarrow{\ell} v_j$.

We call $\overrightarrow{\ell} \in R(G)$ an oriented edge, the edges in $E(G) - R(G)$ are called normal edges, or simply edges. If for all $\ell \in R(G)$ we have an orientation, we say that $\mathfrak{d}$ is an oriented decomposition.

Assumption 2. There is an orientation of $\mathfrak{d}$ such that

(1) For any edge $\ell \in R(G)$ connecting $v_i \in G_{I_i}$ and $v_j \in G_{J_\alpha}$, we have

$$\overrightarrow{\ell} : v_i \xrightarrow{\ell} v_j,$$

and we denote $G_{I_i} \to G_{J_\alpha}$.

(2) For any connected subgraph G_{J_α} and G_{J_β}, all edges between G_{J_α} and G_{J_β} have the same orientation, that is, if there is an edge $\ell \in R(G)$ connecting $v_i \in G_{J_\alpha}$ and $v_j \in G_{J_\beta}$ satisfying $v_i \xrightarrow{\ell} v_j$, then any edge $\ell' \in R(G)$ connecting $v_i' \in G_{J_\alpha}$ and $v_j' \in G_{J_\beta}$ satisfies $v_i' \xrightarrow{\ell} v_j'$. We denote $G_{J_\alpha} \to G_{J_\beta}$.

(3) For any subgraph sequence $G_{J_{\alpha_1}}, \cdots, G_{J_{\alpha_l}}$ satisfying

$$G_{J_{\alpha_1}} \to G_{J_{\alpha_2}} \to \cdots \to G_{J_{\alpha_l}},$$

if there is an edge $\ell \in R(G)$ connecting $v_i \in G_{J_{\alpha_i}}$ and $v_j \in G_{J_{\alpha_j}}$ with $j > i$ then we have

$$v_i \xrightarrow{\ell} v_j, \qquad G_{J_{\alpha_i}} \to G_{J_{\alpha_j}}.$$

(4) For any vertices $v_{j_0}, v_{j_i} \in G$, if there exist a walk of the form

$$v_{j_0}, e_{j_1}, v_{j_1}, e_{j_2}, \cdots, e_{j_i}, v_{j_i},$$

where each edge e_{j_l}, $1 \leq l \leq i$, is the normal edge, then v_{j_0}, v_{j_i} belong to the same subgraph; otherwise v_{j_0}, v_{j_i} belong to different subgraphs.

Let $v_k \xrightarrow{\ell} v_j$, $v_k \in G_{I_i}$ (or G_{J_α}), $v_j \in G_{J_\beta}$. When we consider the subgraphs G_{I_i}, G_{J_β} we attach a half edge ℓ^+ to v_k and attach ℓ^- to v_j.

Let $(V(G), E(G))$ be a graph and $\mathfrak{d}$ be an oriented decomposition satisfying **Assumption 1** and **Assumption 2**. We call the graph G a graph with a oriented decomposition $\mathfrak{d}$, denoted by $(V(G), E(G), \mathfrak{d})$.

Definition 5.9. *Let g, m and ν be nonnegative integers. A $(g, m+\nu, \mathfrak{d})$-weighted dual graph G consists of $(V(G), E(G), \mathfrak{d})$ together with three weights, where*

(1) $(V(G), E(G)$ is a graph, and $\mathfrak{d}$ is an oriented decomposition of $V(G)$,

(2) $\mathfrak{g} : V(G) \to \mathbb{Z}_{\geq 0}$ assigning a nonnegative integer g_v to each vertex v such that

$$g = \sum_{v \in V(G)} g_v + b_1(G)$$

where $b_1(G)$ is the first Betti number of the graph G;

(3) assign m ordered tails $\mathfrak{m} = (t_1, \cdots, t_m)$ to $V(G)$: attach m_v tails to v for each $v \in V(G)$,

(4) assign ν ordered half edges $\mathfrak{l} = (e_1, \cdots, e_\nu)$ to $V(G)$: attach l_v half edges to v for each $v \in V(G)$.

We denote the $(g,m+\nu,\mathfrak{d})$-weighted dual graph G by $(V(G),E(G),\mathfrak{g},\mathfrak{m},\mathfrak{l},\mathfrak{d})$.

We introduce a terminology: a vertex v is called an interior vertex if there is no half edge attached it, it is called boundary vertex if there are some half edges attached it.

Definition 5.10. *Let g, m and ν be nonnegative integers. A H-$(g,m+\nu,\mathfrak{d})$ weighted dual graph G consists of $(V(G),E(G),\mathfrak{g},\mathfrak{m},\mathfrak{l},\mathfrak{d})$ together with three weights:*

(1) $\mathfrak{h}: V(G)\to H_2(M^+,\Re,\mathbb{Z})\bigcup H_2(M^+,\mathbb{Z})$ assigning a $A_v\in H_2(M^+,\Re,\mathbb{Z})$ to each boundary vertex, assigning a $A_v\in H_2(M^+,\mathbb{Z})$ to each interior vertex $v\in\mathfrak{A}$, assigning a 0 to each interior vertex $v\in\mathfrak{B}$. For any G_{I_i} and G_{J_α}, denote

$$A_{I_i}=\sum_{v\in V(G_{I_i})}A_v\in H_2(M^+,\Re,\mathbb{Z}),\qquad A_{J_\alpha}=\sum_{v\in V(G_{J_\alpha})}A_v\in H_2(M^+,\Re,\mathbb{Z}),$$

and

$$A=\sum_{i=1}^{c}A_{I_i}+\sum_{\alpha=1}^{d}A_{J_\alpha}.$$

(2) assign ν ordered weights $\mathbf{k}=(k_1,...,k_\nu)$ and weights $\mathfrak{e}=(\mathfrak{e}_1,\cdots,\mathfrak{e}_\nu)$ to the half edges $\mathfrak{l}=(e_1,\cdots,e_\nu)$ such that $\mathfrak{l}$ becomes weighted half edges $\mathfrak{l}^{(\mathbf{k},\mathfrak{e})}=((k_1,\mathfrak{e}_1)e_1,\cdots,(k_\nu,\mathfrak{e}_\nu)e_\nu)$.

(3) $\mathfrak{k}: R(G)\to\mathbb{Z}^+$, for each $\ell\in R(G)$ with $v_k\xrightarrow{\ell}v_j$, $v_k\in G_{I_i}$ (or G_{J_α}), $v_j\in G_{J_\beta}$, assigning $\mathfrak{k}(\ell)=k_\ell>0$ such that

$$\sum_{e_j\in G_{I_i}}k_j+\sum_{\ell^+\in G_{I_i}}k_\ell>0,\ \text{for any } G_{I_i}$$

and

$$d\lambda(A_{J_\alpha})=\sum_{e_j\in G_{J_\alpha}}k_j\cdot T_{\mathfrak{e}_j}+\sum_{\ell^+\in G_{J_\alpha}}k_{\ell^+}\cdot T_{\ell^+}-\sum_{\ell^-\in G_{J_\alpha}}k_{\ell^-}\cdot T_{\ell^-},\ \text{for any } G_{J_\alpha}.$$

and for each boundary vertex $v\in G_{J_\alpha}$

$$d\lambda(A_v)=\sum_{e_i^+\in\mathfrak{l}_v^+}k_i^+\cdot T_{e_i^+}-\sum_{e_i^-\in\mathfrak{l}_v^-}k_i^-\cdot T_{e_i^-},\tag{86}$$

where $\mathfrak{l}_v^\pm$ is the subset of $\mathfrak{l}^\pm$, the half edges attached to v

We denote the H-$(g,m+\nu,\mathfrak{d})$ weighted dual graph G by $(V(G),E(G),\mathfrak{g},\mathfrak{m},\mathfrak{l}^{(\mathbf{k},\mathfrak{e})},\mathfrak{k},\mathfrak{h},\mathfrak{d})$.

By a leg of G we mean either a tail or a half-edge.

Definition 5.11. *Let G be a $(V(G),E(G),\mathfrak{g},\mathfrak{m},\mathfrak{l}^{(\mathbf{k},\mathfrak{e})},\mathfrak{k},\mathfrak{h},\mathfrak{d})$ graph. A vertex v is called stable if one of the following holds:*

(1) $2g_v+val(v)\geq 3$, where $val(v)$ denotes the sum of the number of legs attached to v;

(2) $A_v\neq 0$, when $v\in\mathfrak{A}$;

(3) $d\lambda(A_v)\neq 0$ when $v\in\mathfrak{B}$.

G is called stable if all vertices are stable.

Let G be a stable $(V(G), E(G), \mathfrak{g}, \mathfrak{m}, \mathfrak{l}^{(\mathbf{k},\mathfrak{e})}, \mathfrak{k}, \mathfrak{h}, \mathfrak{d})$ graph. Then all subgraphs G_{I_i}, G_{J_α} are stable.

Two H-$(g, m+\nu, \mathfrak{d})$ weighted dual graphs G_1 and G_2 are called isomorphic if there exists a bijection T between their vertices and edges keeping oriented decomposition and all weights.

Let $S_{\mathfrak{g},\mathfrak{m},\mathfrak{l}^{(\mathbf{k},\mathfrak{e})},\mathfrak{k},\mathfrak{h},\mathfrak{d}}$ be the set of isomorphic classes of H-$(g, m+\nu, \mathfrak{d})$ weighted dual graphs. Given g, m, ν, $A \in H_2(M^+, \Re, \mathbb{Z})$ and two weights $\mathbf{k} = (k_1, ..., k_\nu)$, $\mathfrak{e} = (\mathfrak{e}_1, \cdots, \mathfrak{e}_\nu)$, denote by $S_{g,m+\nu,A,\mathfrak{l}^{(\mathbf{k},\mathfrak{e})}}$ the union of all possible $S_{\mathfrak{g},\mathfrak{m},\mathfrak{l}^{(\mathbf{k},\mathfrak{e})},\mathfrak{k},\mathfrak{h},\mathfrak{d}}$.

5.6 Li-Ruan's Compactification

In this section we state Li-Ruan's compactification on the moduli space of maps in terms of language of graphs.

Let G be a stable $(V(G), E(G), \mathfrak{g}, \mathfrak{m}, \mathfrak{l}^{(\mathbf{k},\mathfrak{e})}, \mathfrak{k}, \mathfrak{h}, \mathfrak{d})$ graph with N vertices $(v_1, ..., v_N)$, m tails and ν half edges, and $(\Sigma, \mathbf{y}, \mathbf{p})$ be a semi-stable curve with m marked points and ν puncture points. Let $A \in H_2(M^+, \Re, \mathbb{Z})$. A stable J-holomorphic map of type G is a quadruple

$$(u; \Sigma, \mathbf{y}, \mathbf{p})$$

where $u : \Sigma \to (M^+)'$ is a continuous map, $(M^+)'$ is obtained by attaching some $\mathbb{R} \times \widetilde{M}$ to M^+, satisfying the following conditions:

[A-1] $\Sigma = \bigcup_{v=1}^{N} \Sigma_v$, where each $v \in V(G)$ represents a smooth component Σ_v of Σ.

[A-2] for the i-th tail attached to the vertex v there exists the i-th marked point $y_i \in \Sigma_v$, m_v is equal to the number of the marked points on Σ_v,

[A-3] for the j-th half edge attached to the vertex v there exists j-th puncture point $p_j \in \Sigma_v$, l_v is equal to the number of puncture points on Σ_v;

[A-4] if there is an edge connected the vertices v and w, then there exists a node between Σ_v and Σ_w, the number of edges between v and w is equal to the number of node points between Σ_v and Σ_w;

[A-6] the restriction of u to each component Σ_v is J-holomorphic.

[A-7] u converges exponentially to $(k_1 \cdot T_{\mathfrak{e}_1}, \cdots, k_\nu \cdot T_{\mathfrak{e}_\nu})$ periodic orbits $(x_{k_1}, ..., x_{k_\nu})$ satisfying $x_{k_i} \subset \mathcal{F}_{\mathfrak{e}_i}$ as the variable tends to the puncture $(p_1, ..., p_\nu)$; more precisely, u satisfies (27)-(29);

[A-8] let q be a nodal point of Σ. Suppose q is the intersection point of Σ_v and Σ_w associated to the edge $\ell \in R(G)$. Then $k_\ell > 0$, $u|_{\Sigma_v}$ and $u|_{\Sigma_w}$ converge exponentially to the same k_ℓ periodic orbit x_{k_ℓ} on $\widetilde{M}$ as the variables tend to the nodal point q.

[A-9] For any $v \in V(G)$, $[u(\Sigma_v)] = A_v \in H_2(M^+, \Re, \mathbb{Z})$ when v is a boundary vertex; $[u(\Sigma_v)] = A_v \in H_2(M^+, \mathbb{Z})$ when $v \in \mathfrak{A}$ is an interior vertex, $[u(\Sigma_v)] = 0$ when $v \in \mathfrak{B}$ is an interior vertex; $A = \sum_{i=1}^{v} A_v$.

Definition 5.12. *Two stable J-holomorphic maps $\Gamma = (u, (\Sigma, \mathbf{j}), \mathbf{y}, \mathbf{p})$ and $\check{\Gamma} = (\check{u}, (\check{\Sigma}, \check{\mathbf{j}}), \check{\mathbf{y}}, \check{\mathbf{p}})$ of type G are called equivalent if there exists a diffeomorphism $\varphi : \Sigma \to \check{\Sigma}$ such that it can be lifted to bi-holomorphic isomorphisms $\varphi_{vw} : (\Sigma_v, j_v) \to (\check{\Sigma}_w, \check{j}_w)$ for each component Σ_v of Σ, and*

- **(1)** *$\varphi(y_i) = \check{y}_i$, $\varphi(p_j) = \check{p}_j$ for any $1 \leq i \leq l$, $1 \leq j \leq \nu$,*
- **(2)** *$u(\Sigma_v)$ and $\check{u} \circ \varphi(\Sigma_v)$ lie in the same holomorphic block for any Σ_v,*
- **(3)** *In case $v \in \mathfrak{A}$ we have $\check{u} \circ \varphi = u$ on Σ_v; In case $v \in \mathfrak{B}$, we have*

  $$\check{\tilde{u}} \circ \varphi = \tilde{u}, \;\; \check{a} \circ \varphi = a + C \;\text{ on } \Sigma_v.$$

  Moreover, near every periodic orbit x, $\tilde{u}$ and $\check{\tilde{u}} \circ \varphi$ may differ by a canonical coordinate transformation (26).

Denote by $\mathcal{M}_G$ the space of the equivalence class of stable J-holomorphic maps of type G.

Remark 5.13. *Every subgraph G_{I_i} determines a holomorphic block of type G_{I_i} in M^+, every subgraph G_{J_α} determines a holomorphic rubber block of type G_{J_α} in $\mathbb{R} \times \widetilde{M}$. Roughly speaking, $\mathcal{M}_G$ is the gluing of several holomorphic blocks.*

We define the automorphism group of u:

$$\begin{aligned} Aut(u) = \{\varphi \mid \varphi : \Sigma \to \Sigma \text{ is a holomorphic isomorphic such that } (\mathbf{1}), (\mathbf{2}) \\ \text{and } \mathbf{(3)} \text{ hold in } \mathbf{Definition}\ 5.12.\} \end{aligned}$$

It is easy to see

Lemma 5.14. *For any stable holomorphic map u of type G the automorphism group $Aut(u)$ is finite .*

Given g, m, ν, $A \in H_2(M^+, \Re, \mathbb{Z})$, two weights $\mathbf{k} = (k_1, ..., k_\nu)$ and $\mathfrak{e} = (\mathfrak{e}_1, \cdots, \mathfrak{e}_\nu)$, we define

$$\overline{\mathcal{M}}_A(M^+, C; g, m + \nu, (\mathbf{k}, \mathfrak{e})) = \bigcup_{G \in S_{g, m+\nu, A, \mathfrak{l}(\mathbf{k}, \mathfrak{e})}} \mathcal{M}_G.$$

This gives a stratification of $\overline{\mathcal{M}}_A(M^+, C; g, m + \nu, (\mathbf{k}, \mathfrak{e}))$. Denote by $\mathcal{D}^{J,A}_{g, m+\nu, \mathbf{k}, \mathfrak{e}}$ the number of all possible $S_{\mathfrak{g}, \mathfrak{m}, \mathfrak{l}(\mathbf{k}, \mathfrak{e}), \mathfrak{k}, \mathfrak{h}, \mathfrak{d}}$. The following Lemma is obvious.

Lemma 5.15. *$\mathcal{D}^{J,A}_{g, m+\nu, \mathbf{k}, \mathfrak{e}}$ is finite.*

We immediately obtains

Theorem 5.16. *$\overline{\mathcal{M}}_A(M^+, C; g, m + \nu, (\mathbf{k}, \mathfrak{e}))$ is compact.*

Remark 5.17. *Let $u : S^2 \longrightarrow \mathbb{R} \times \tilde{M}$ be a J-holomorphic map with $\tilde{E}(u) = 0$. Then u must be $u = (kTs + d, x(kt + \vartheta_0))$. If the number of the special points (including nodal points, puncture points and marked points) ≤ 2, u is called a unstable second class ghost bubble. In our compactification there is no unstable second class ghost bubble.*

6 Gluing theory-Pregluing

The gluing is the inverse of the degeneration. In section §5 we see that for a sequence $\Gamma^{(i)} = (u^{(i)}, \Sigma^{(i)}; \mathbf{y}^{(i)}, \mathbf{p}^{(i)}) \in \mathcal{M}_A(M^+, C; g, m+\nu, (\mathbf{k}, \mathfrak{e}))$, each component degenerates independently, and the bubble trees are also constructed independently, furthermore each node of a connected component degenerates also independently. So we also glue independently for each nodal points. Recall that there are some freedoms of choosing the coordinates a, ϑ for M^+ and $\mathbb{R} \times \widetilde{M}$ (see subsection §2.3).

For any r we glue M^+ and $\mathbb{R} \times \widetilde{M}$ with parameter r to get again M^+. We cut off the part of M^+ and $\mathbb{R} \times \widetilde{M}$ with cylindrical coordinate $|a^{\pm}| > \frac{3lr}{2}$ and glue the remainders along the collars of length lr of the cylinders with the gluing formulas:

$$a_1 = a_2 + 2lr. \tag{87}$$

6.1 Gluing one relative node for $M^+ \cup (\mathbb{R} \times \widetilde{M})$

Let $b = (u_1, u_2; \Sigma_1 \wedge \Sigma_2, j_1, j_2) \in \mathcal{M}_{A_1}(M^+, C; g_1, m_1 + 1, (k, \mathfrak{e}_1)) \times \mathcal{M}_{A_2}(\mathbb{R} \times \widetilde{M}, g_2, m_2 + \nu + 1, (k, \mathfrak{e}_1), (\mathbf{k}^+, \mathfrak{e}^+))$, where (Σ_1, j_1) and (Σ_2, j_2) are smooth Riemann surfaces of genus g_1 and g_2 joining at q and $u_1 : \Sigma_1 \to M^+$, $u_2 : \Sigma_2 \to \mathbb{R} \times \widetilde{M}$ are J-holomorphic maps such that $u_i(z)$ converge to the same kT-periodic orbit x as $z \to q$. To describe the maps u_i we choose a local pseudo-Darboux coordinate system $(a_i, \vartheta_i, \mathbf{w})$, $i = 1, 2$, near x. Suppose that

$$a_i(s_i, t_i) - kTs_i - \ell_i \to 0 \quad \vartheta_i(s_i, t_i) - kt_i - \vartheta_{i0} \to 0.$$

If we choose a different origin in the periodic orbit x, we have a different coordinate system ϑ_i^*. Suppose that

$$\vartheta_i^* = \vartheta_i + \tau_i.$$

Obviously,

$$\vartheta_i^*(s_i, t_i) - kt_i - (\vartheta_{i0} + \tau_i) \to 0.$$

Without loss of generality we assume that $\vartheta_{10} = 0$. Set $\tau = -\vartheta_{20}$. We consider τ to be a parameter satisfying $0 \le \tau < 1$. Then

$$\vartheta_1 = \vartheta_2 + \tau \tag{88}$$

Given gluing parameters $(r) = (r, \tau)$ we construct a surface $\Sigma_{(r)} = \Sigma_1 \#_{(r)} \Sigma_2$ with gluing formulas:

$$s_1 = s_2 + \tfrac{2lr}{Tk} \tag{89}$$

$$t_1 = t_2 + \tfrac{\tau + n}{k} \tag{90}$$

for some $n \in Z_k$.

To get a pregluing map $u_{(r)}$ from $\Sigma_{(r)}$ we set

$$u_{(r)} = \begin{cases} u_1 \quad on \;\; \Sigma_{10} \bigcup \left\{(s_1, t_1) | 0 \le s_1 \le \frac{lr}{2Tk}, t_1 \in S^1\right\} \\ (kTs_1, x(kTt_1)) \quad\;\; on \;\; \left\{(s_1, t_1) | \frac{3lr}{4Tk} \le s_1 \le \frac{5lr}{4Tk}, t_1 \in S^1\right\} \\ u_2 \quad on \;\; \Sigma_{20} \bigcup \left\{(s_2, t_2) | 0 \ge s_2 \ge -\frac{lr}{2Tk}, t_2 \in S^1\right\}, \end{cases}$$

where $x(kTt_1) = (kt_1, 0)$ as $\vartheta_{10} = 0$.

To define the map $u_{(r)}$ in the remaining part we fix a smooth cutoff function $\beta : \mathbb{R} \to [0,1]$ such that

$$\beta(s) = \begin{cases} 1 & if \;\; s \geq 1 \\ 0 & if \;\; s \leq 0 \end{cases}$$

and $\sqrt{1-\beta}$ is a smooth function, $|\beta'(s)| \leq 2$. We assume that r is large enough such that u_i maps the tube $\{(s_i,t_i) | |s_i| \geq \frac{lr}{4Tk}, t_i \in S^1\}$ into a domain with pseudo-Darboux coordinates $(a_i, \vartheta_i, \mathbf{w})$. We write $u_{(r)} = \big(a_{(r)}, \widetilde{u}_{(r)}\big)$ and define

$$a_{(r)} = kTs_1 + \Big(\beta(3 - \tfrac{4Tks_1}{lr})(a_1(s_1,t_1) - kTs_1) + \beta(\tfrac{4Tks_1}{lr} - 5)(a_2(s_2,t_2) - kTs_2)\Big), \tag{91}$$

$$\widetilde{u}_{(r)} = x(kTt_1) + \Big(\beta(3 - \tfrac{4Tks_1}{lr})(\widetilde{u}_1(s_1,t_1) - x(kTt_1)) + \beta(\tfrac{4Tks_1}{lr} - 5)(\widetilde{u}_2(s_2,t_2) - x(kTt_2))\Big), \tag{92}$$

where $x(kTt_2) = (kt_2, 0)$ as (90) and (88). It is easy to check that $u_{(r)}$ is a smooth function.

6.2 Gluing two relative nodal points for $M^+ \cup (\mathbb{R} \times \widetilde{M})$

Recall Lemma 5.8, we glue two relative nodal points independently.

Let $b = (u_1, u_2; \Sigma_1 \wedge \Sigma_2, j_1, j_2) \in \mathcal{M}_{A_1}(M^+, C; g_1, m_1 + 2, (\mathbf{k}, \mathfrak{e})) \times \mathcal{M}_{A_2}(\mathbb{R} \times \widetilde{M}, C; g_2, m_2 + 2, (\mathbf{k}^-, \mathfrak{e}^-), (k', \mathfrak{e}'))$, where $\mathbf{k} = \mathbf{k}^- = (k_1, k_2), \mathfrak{e} = \mathfrak{e}^- = (\mathfrak{e}_1, \mathfrak{e}_2)$, and k' satisfying $\sum_{j=1}^{2} k_j < k'$. Here (Σ_1, j_1) and (Σ_2, j_2) are smooth Riemann surfaces of genus g_1 and g_2 joining at q_1, q_2, and $u_1 : \Sigma_1 \to M^+$, $u_2 : \Sigma_2 \to \mathbb{R} \times \widetilde{M}$ are J-holomorphic maps such that $u_j(z)$ converge to the same $k_jT_{\mathfrak{e}_j}$-periodic orbit $x_j(k_jT_{\mathfrak{e}_j})$ as $z \to q_j, j = 1, 2$ (see Figure 2). Choose cylinder coordinates (s_{1j}, t_{1j}) and (s_{2j}, t_{2j}) on Σ_1 and Σ_2 near node q_j. We choose local pseudo- Darboux coordinate systems $(a_1, \vartheta_{1j}, \mathbf{w}_j)$ on the cylinder end of M^+, $(a_2, \vartheta_{2j}, \mathbf{w}_j)$ on $\mathbb{R} \times \widetilde{M}$ near x_j, where $\mathbf{w}_j$ is a local coordinates near x_j. Suppose that

$$a_1(s_{1j}, t_{1j}) - kTs_{1j} - \ell_{1j} \to 0, \quad \vartheta_{1j}(s_{1j}, t_{1j}) - kt_{1j} - \vartheta_{1j0} \to 0, \quad j = 1, 2,$$

$$a_2(s_{2j}, t_{2j}) - kTs_{2j} - \ell_{2j} \to 0, \quad \vartheta_{2j}(s_{2j}, t_{2j}) - kt_{2j} - \vartheta_{2j0} \to 0, \quad j = 1, 2.$$



Figure 2:

Without loss of generality we assume that $\ell_{1j} = \ell_{2j} = 0$, for $j = 1, 2$. Denote $T_j = T_{\mathfrak{e}_j}$, $j = 1, 2$.

For any parameter $\varrho > 0$, we can gluing M^+ and $\mathbb{R}\times\widetilde{M}$ to get M^+_ϱ with gluing formula:

$$a_1 = a_2 + 2l\varrho. \tag{93}$$

For each x_j we take $\mathcal{T}$ translation:

$$a_{1j} = a_1 - c_{1j}, \;\; a_{2j} = a_2 - c_{2j} \tag{94}$$

for some constants $c_{1j} \geq 0$. Then $(a_{1j}, \vartheta_{1j}, \mathbf{w}_j)$ and $(a_{2j}, \vartheta_{2j}, \mathbf{w}_j)$ are local coordinate systems near x_j over cylindrical end of M^+ and over $\mathbb{R}\times\widetilde{M}$.

Choose R_0 such that

$$\sum_{i=1}^{2}\sum_{j=1}^{2}\widetilde{E}(u_i;|s_{ij}|\geq \frac{R_0}{2}) \leq \frac{\min\{\hbar, T\}}{8}, \qquad 2\mathcal{C}_1\mathfrak{c}_1^{-1}e^{-\frac{\mathfrak{c}R_0}{4}} \leq \frac{\hbar_1}{8}, \tag{95}$$

where $\hbar, \mathcal{C}_1, \mathfrak{c}$ and $\hbar_1$ are the constants in Theorem 2.9 and Theorem 2.10.

For any $r_j \geq R_0$ we glue M^+ and $\mathbb{R}\times\widetilde{M}$ in coordinates a_{1j} and a_{2j} with the gluing formula:

$$a_{1j} = a_{2j} + 2lr_j. \tag{96}$$

Now we return to the coordinates $a_i, i = 1, 2$. By relation (94) the gluing formula can be re-written as

$$a_1 = a_2 + c_{1j} - c_{2j} + 2lr_j. \tag{97}$$

Choose

$$c_{1j} = -c_{2j} = l(\rho - r_j). \tag{98}$$

By (98), in order to get M^+_ϱ we need

$$R_0 \leq r_j \leq \varrho, \qquad j = 1, 2. \tag{99}$$

Now we glue J-holomorphic maps. We express (u_1, u_2) in terms of the coordinates (a_{1j}, ϑ_{1j}) and (a_{2j}, ϑ_{2j}):

$$a_{1j}(s_{1j}, t_{1j}) - k_jT_js_{1j} + c_{1j} \to 0, \quad \vartheta_{1j}(s_{1j}, t_{1j}) - k_jt_{1j} - \vartheta_{1j0} \to 0, \qquad j = 1, 2,$$

$$a_{2j}(s_{2j}, t_{2j}) - k_jT_js_{2j} + c_{2j} \to 0, \quad \vartheta_{2j}(s_{2j}, t_{2j}) - k_jt_{2j} - \vartheta_{2j0} \to 0, \qquad j = 1, 2.$$

As in subsection §6.1 we assume $\vartheta_{1j0} = 0$ and consider $\tau_j = -\vartheta_{2j0}$ as parameters, and construct a surface $\Sigma_{(r_j)} = \Sigma_1\#_{(r_j)}\Sigma_2$ with gluing formulas:

$$s_{1j} = s_{2j} + \tfrac{2lr_j}{k_jT_j}, \qquad j = 1, 2, \tag{100}$$

$$t_{1j} = t_{2j} + \tfrac{\tau_j + n_j}{k_j}, \qquad j = 1, 2, \tag{101}$$

for some $n_j \in Z_{k_j}, j = 1, 2$. By (98), in terms of ϱ the gluing formulas (100) and (101) can be written as

$$s_{1j} = s_{2j} + \tfrac{2l\rho - c_{1j} + c_{2j}}{k_jT_j}, \qquad j = 1, 2, \tag{102}$$

$$t_{1j} = t_{2j} + \tfrac{\tau_j + n_j}{k_j}, \qquad j = 1, 2, \tag{103}$$

for some $n_j \in Z_{k_j}, j = 1, 2$. We construct pre-gluing map $u_{(r_j)}$ as in subsection §6.1. Then we have

$$u_{(r_1)} = u_{(r_2)} = u_1, \text{ in } \bigcap_{j=1}^{2} \{|s_{1j}| \leq \frac{lr_j}{4T_j k_j}\}, \qquad u_{(r_1)} = u_{(r_2)} = u_2, \text{ in } \bigcap_{j=1}^{2} \{|s_{2j}| \leq \frac{lr_j}{4T_j k_j}\}.$$

The pre-gluing above can be generalized immediately to the case of gluing several nodal points.

6.3 Norms on $C^\infty(\Sigma_{(r)}; u^*_{(r)} TM^+_{(r)})$

We only consider the case of gluing one node, the other cases are the same. For any $\eta \in C^\infty(\Sigma_{(r)}; u^*_{(r)} TM^+_{(r)} \otimes \wedge^{0,1}))$, let η_i be its restriction to the part $\Sigma_{i0} \bigcup \{(s_i, t_i) |\ |s_i| < \frac{3lr}{2kT}\}$, extended by zero to yield a section over Σ_i. Define

$$\|\eta\|_{p,\alpha,r} = \|\eta_1\|_{\Sigma_1,p,\alpha} + \|\eta_2\|_{\Sigma_2,p,\alpha}. \tag{104}$$

Denote the resulting completed spaces by $L^{p,\alpha}_r$.

We define a norm $\|\cdot\|_{1,p,\alpha,r}$ on $C^\infty(\Sigma_{(r)}; u^*_{(r)} TM^+_{(r)})$. For any section $h \in C^\infty(\Sigma_{(r)}; u^*_{(r)} TM^+_{(r)})$, denote

$$h_0 = \int_{\mathbb{S}^1} h\left(\frac{lr}{Tk}, t\right) dt, \tag{105}$$

$$h_1 = (h - \hat{h}_0)\beta\left(\frac{3}{2} - \frac{Tks_1}{lr}\right) \tag{106}$$

$$h_2 = (h - \hat{h}_0)\left[1 - \beta\left(\frac{3}{2} - \frac{Tks_1}{lr}\right)\right]. \tag{107}$$

We define

$$\|h\|_{1,p,\alpha,r} = \|h_1\|_{\Sigma_1,1,p,\alpha} + \|h_2\|_{\Sigma_2,1,p,\alpha} + |h_0|. \tag{108}$$

Denote the resulting completed spaces by $\mathcal{W}^{1,p,\alpha}_r$.

We introduce some notations. Choose the cylinder coordinates near puncture points and nodal points. Denote

$$\mathfrak{D}_i(R_0) = \Sigma_{i0} \cup \{(s_i, t_i) |\ |s_i| \leq R_0\}, \ \ i = 1, \ 2,$$
$$\mathfrak{D}(R_0) = \mathfrak{D}_1(R_0) \bigcup \mathfrak{D}_2(R_0).$$

Choose R_0 such that

$$\sum_{i=1}^{2} \widetilde{E}(u_i; |s_i| \geq \frac{R_0}{2}) \leq \frac{\min\{\hbar, T\}}{8}, \qquad 4\mathcal{C}_1 \mathfrak{c}_1^{-1} e^{-\frac{\mathfrak{c}R_0}{4}} \leq \frac{\hbar_1}{8}, \tag{109}$$

We let $\frac{lr}{Tk} >> 4R_0$.

Lemma 6.1. *There exists $\epsilon > 0$ such that for any J-holomorphic map $v : \Sigma_{(r)} \to M^+_{(r)}$ with $v = \exp_{u_{(r)}}(h)$, if*

$$|h\ |_{\mathfrak{D}_i(R_0)}\ |_{C^1} \leq \epsilon \ \ for \ \ i = 1, 2, \tag{110}$$

then for any $0 < \alpha < \mathfrak{c}$

$$\|h\|_{1,p,\alpha,r} \leq \frac{\hbar_1}{4},$$

where $\mathfrak{c}$ is the constant in Theorem 2.9.

Proof. Note that

$$\widetilde{E}(v;|s_i|\geq R_0/2)=\int_{R_0/2\leq s_1\leq \frac{2lr}{Tk}-R_0/2} v^*d\lambda=\int_{v(\frac{2lr}{Tk}-R_0/2,S^1)}\lambda-\int_{v(R_0/2,S^1)}\lambda.$$

Denote $s_0=R_0/2$. A direct calculation gives us

$$\begin{aligned}&\left|\int_{v(s_0,S^1)}\lambda-\int_{u_1(s_0,S^1)}\lambda\right|=\left|\int_{S^1}\lambda(v_t)(s_0,t)dt-\int_{S^1}\lambda((u_1)_t)(s_0,t)dt\right|\\ &\leq\left|\int_{S^1}\lambda(P_{u_1,v}(u_1)_t))(s_0,t)dt-\int_{S^1}\lambda((u_1)_t)(s_0,t)dt\right|+\left|\int_{S^1}\lambda(d\exp_{u_1}h_t)(s_0,t)dt\right|\\ &\leq C|h\,|_{\mathfrak{D}_1(R_0)}\,|_{C^1} \qquad (111)\end{aligned}$$

Then

$$\widetilde{E}(v;|s_i|\geq R_0/2)\leq\widetilde{E}(u_i;|s_i|\geq R_0/2)+C\sum|h\,|_{\mathfrak{D}_i(R_0)}\,|_{C^1}.$$

It follows that

$$\widetilde{E}(v;\Sigma_{(r)}-\mathfrak{D}(R_0/2))\leq\frac{\min\{\hbar,T\}}{4},$$

when $0<\epsilon<\frac{\min\{\hbar,T\}}{8C}$. Then by Theorem 2.9 we have

$$|\nabla v\,|_{\Sigma_{(r)}-\mathfrak{D}(R_0)}\,|\leq\mathcal{C}_1e^{-\mathfrak{c}R_0/2} \qquad (112)$$

for all $R_0\leq s_1\leq\frac{lr}{Tk}-R_0$. Together with (110) we have

$$\|h\,|_{\mathfrak{D}(R_0)}\,\|_{1,p,\alpha}\leq\frac{\hbar_1}{8}$$

when ϵ small enough. Then lemma follows. $\Box$

The lemma can be generalized to the case of gluing several Riemann surfaces and several nodal points.

7 Gluing theory–Regularization

7.1 Local regularization

We fist discuss the top strata, then consider lower stratas.

7.1.1 Top strata.

Let $b=(u,(\Sigma,j);\mathbf{y},\mathbf{p})\in\overline{\mathcal{M}}_A(M^+,C;g,m+\nu,(\mathbf{k},\mathfrak{e}))$. Here Σ is a smooth Riemann surface of genus g, j is a complex structure (including marked points), which is standard near each puncture point. We introduce the holomorphic cylindrical coordinates on Σ near each puncture points p_i.

We discuss two different cases:

(1.1) Σ is stable.

Denote by O_j a neighborhood of complex structures on (Σ, j). Note that we change the complex structure in the compact set $\mathfrak{D}(R_0)$ of Σ away from the puncture points. A neighborhood $\mathcal{U}_b$ of b can be described as

$$O_j \times \{\exp_u(h + \hat{h}_0); h \in \mathcal{C}(\Sigma; u^*TM^+), h_0 \in \mathbb{H}, \|h\|_{1,p,\alpha} + |h_0| < \epsilon\}/stab_b,$$

where $\mathbb{H} = \bigoplus_{j=1}^{\nu}(T_{p_j}(\mathcal{F}_{\mathfrak{e}_j}) \oplus (span\{\frac{\partial}{\partial a}\}))$.

(1.2) Σ is unstable.

In this case the automorphism group Aut_Σ is infinite. One must construct a slice of the action Aut_Σ and construct a neighborhood $\mathcal{U}_b$ of b. This is done in [7]. We omit it here.

There is a neighborhood U of b such that $\mathcal{E}$ is trivialized over U, more precisely , $P(u, v)$ gives a isomorphism $\mathcal{E}|_u \to \mathcal{E}|_v$ for any $v \in U$. Choose $K_b \subset \mathcal{E}|_u$ to be a finite dimensional subspace such that

$$K_b + imageD_u = \mathcal{E}|_u, \tag{113}$$

where $D_u : W^{1,p,\alpha}(u^*TM^+) \to \mathcal{E}|_u$. Without loss of generality we may assume that every element of K_b is smooth along u and supports in the common compact subset $\mathfrak{D}(R_0)$. Define a thickned Fredholm system $(K_b \times U, K_b \times \mathcal{E}|_U, S_b)$, where

$$\mathcal{S}(\kappa, v) = \bar{\partial}_J v + P_{u;v}\kappa \in \mathcal{E}_v, \quad \forall\ (\kappa, v) \in K_b \times U.$$

By (113), there exists a smaller neighborhood of b, still denoted by U, such that $D\mathcal{S}_{(\kappa,v)} : K_b \times W^{1,p,\alpha} \to L^{p,\alpha}$ is surjective for any $(\kappa, v) \in K_b \times U$. Then for each $b' = (\kappa, v) \in K_b \times U$ there exists a right inverse

$$Q_{b'} : L^{p,\alpha} \to K_b \times W^{1,p,\alpha}$$

such that $\|Q_{b'}\| \leq C_1$. Obviously, $D\mathcal{S}_{b'} : K_b \times \mathcal{W}^{1,p,\alpha} \to L^{p,\alpha}$ is also surjective, $Q_{b'} : L^{p,\alpha} \to K_b \times \mathcal{W}^{1,p,\alpha}$ is also a right inverse of $D\mathcal{S}_{b'}$.

A pair $(\kappa, v) \in K_b \times U$ is called a perturbed J-holomorphic map if

$$\bar{\partial}_J v + P_{u;v}\kappa = 0.$$

7.1.2 Lower strata.

We shall consider two cases for simplicity, the discussion for general cases are the same. Let D be a strata whose domain has two components (Σ_1, j_1) and (Σ_2, j_2) joining at p.

Case 1. Let $b = (u_1, u_2; \Sigma_1 \wedge \Sigma_2, j_1, j_2)$, where (Σ_1, j_1) and (Σ_2, j_2) are smooth Riemann surfaces of genus g_1 and g_2 joining at p and $u_i : \Sigma_i \to M^+$ are J-holomorphic maps with $u_1(p) = u_2(p)$. Suppose that both (u_i, Σ_i, j_i) are stable. A neighborhood $\mathcal{U}_b^D$ of b in the strata D can be described as

$$O_{j_1} \times O_{j_2} \times \Big\{ \Big(\exp_{u_1}(h_1 + \hat{h}_{10}), \exp_{u_2}(h_2 + \hat{h}_{20})\Big) |h_i \in W^{1,p,\alpha}(\Sigma; {u_i}^*TM^+) \,,$$
$$h_{10} = h_{20} \in T_{u_1(p)}M^+, \|h_i\|_{1,p,\alpha} + |h_0| < \epsilon \Big\} /stab_u.$$

The fiber of $\mathcal{E}_D(M^+, g, m+\nu) \to \mathcal{B}_D(M^+, g, m+\nu)$ at b is $L^{p,\alpha}(u_1^*TM^+ \otimes \wedge^{0,1}) \times L^{p,\alpha}(u_2^*TM^+ \otimes \wedge^{0,1})$.

Case 2. Let $b = (u_1, u_2; \Sigma_1 \wedge \Sigma_2, j_1, j_2)$, where (Σ_1, j_1) and (Σ_2, j_2) are smooth Riemann surfaces of genus g_1 and g_2 joining at q, $u_1 : \Sigma_1 \to M^+$ and $u_2 : \Sigma_2 \to \mathbb{R} \times \widetilde{M}$ are J-holomorphic maps such that u_1 and u_2 converge to a same kT-periodic orbit $x(kTt) \subset \mathcal{F}_j$ for some j when z_1 and z_2 converge to q. Suppose that both (u_i, Σ_i, j_i) are stable.

We first describe the neighborhood of $u_2 : \Sigma_2 \to \mathbb{R} \times \widetilde{M}$ in the space of $\mathcal{W}^{1,p,\alpha}$-maps. The R-action on $\mathbb{R}\times\widetilde{M}$ and S^1-action on every periodic orbit naturally induce the actions on $W^{1,p,\alpha}(\Sigma_2, \mathbb{R}\times \widetilde{M})$, which we need mod. Write $u(s,t) = (a(s,t), \tilde{u})$. For any C, the $\mathbb{R}$-action is defined as

$$\mathbb{R}_C \circ u(s,t) = (C + a(s,t), \tilde{u}).$$

We construct a local R-action slice as follows. We fix a point $p \in \Sigma_2$ and choose coordinates a such that the $u_2(p) = (0, \star)$. Let h be a vector field along u_2. We write

$$h = b_1 \frac{\partial}{\partial a} + \tilde{h},$$

where $\tilde{h} \in W^{1,p,\alpha}(\tilde{u}_2^*(T\widetilde{M}))$ is a vector field along $\tilde{u}_2$. Then $h = b_1 \frac{\partial}{\partial a} + \tilde{h}$ can be considered in a natural way as a $\mathbb{R}$ equivalent vector field along $\mathbb{R} \circ u$. We can construct a equivalent tubular neighborhood around the orbit $\mathbb{R} \circ u$. The point-wise exponential map give us a Banach manifold structure on the space of $\mathbb{R}$ equivalence class of $W^{1,p,\alpha}(\Sigma_2, \mathbb{R} \times \widetilde{M})$. To simplify notations we will use $W^{1,p,\alpha}(\Sigma_2, \mathbb{R} \times \widetilde{M})$ to denote the space of $\mathbb{R}$ equivalence class if no danger of confusion.

On the other hand, we need mod S^1-action on every periodic orbit. Let x be a periodic orbit. We choose a local pseudo-Darboux coordinate system $(a, \vartheta, \mathbf{w})$, near x. If we choose another origin in S^1 we get another local pseudo-Darboux coordinate system $(a', \vartheta', \mathbf{w})$, which differ by a canonical coordinates transformation (26). The vector $h = b_1 \frac{\partial}{\partial a} + \tilde{h}$ is independent of the local pseudo-Darboux coordinates, so independent of the choice of the origin.

For simplicity we assume that $\Sigma_i, i = 1, 2$ has no puncture points. A neighborhood $\mathcal{U}_b^D$ of b in D can be described as

$$O_{j_1} \times O_{j_2} \times \Big\{ (\exp_{u_1}(h_1 + \hat{h}_{10}), \exp_{u_2}(h_2 + \hat{h}_{20})) | h_1 \in W^{1,p,\alpha}(\Sigma; {u_1}^* TM^+),$$

$$h_2 \in W^{1,p,\alpha}(\Sigma; {u_2}^* T(\mathbb{R} \times \widetilde{M})), h_{10} = h_{20} \in \mathbb{H}, \quad \Sigma_{i=1}^2 \|h_i\|_{1,p,\alpha} + |h_{10}| < \epsilon \Big\} / stab_u,$$

where $\mathbb{H} = T_q(\mathcal{F}_j) \oplus (span\{\frac{\partial}{\partial a}\})$. The fiber of $\mathcal{E}_D \to \mathcal{B}_D$ at b is $L^{p,\alpha}(u_1^* TM^+ \otimes \wedge^{0,1}) \times L^{p,\alpha}(u_2^*(\mathbb{R} \times \widetilde{M}) \otimes \wedge^{0,1})$.

We use the gluing argument to describe the neighborhoods $\mathcal{U}_b$ of b in $\mathcal{B}_A(M^+, C; g, m+\nu, (\mathbf{k}, \mathfrak{e}))$. Denote $Glu_{T_0} = \{(r, \tau) | 0 \leq \tau < 1, T_0 \leq r \leq \infty\}$. Then

$$\mathcal{U}_b = \bigcup_{T_0 \leq r < \infty} B(u_{(r)}, \epsilon)$$

where $B(u_{(r)}, \epsilon) = \{u \in \mathcal{B} | \; u = \exp_{u_{(r)}}(h + \hat{h}_0), \quad \|h\|_{1,p,\alpha} + |h_0| < \epsilon\}$.

Choose $K_b = (K_1, K_2) \subset \mathcal{E}|_b = (\mathcal{E}_{u_1}, \mathcal{E}_{u_2})$ to be a finite dimensional subspace such that

$$K_1 + imageD_{u_1} = \mathcal{E}_{u_1}, \quad K_2 + imageD_{u_2} = \mathcal{E}_{u_2}.$$

where $D_{u_i} : W^{1,p,\alpha} \to \mathcal{E}_{u_i}$. We may assume that every element of K_i is smooth along u_i and supports in the compact subset $\mathfrak{D}(R_0)$ of Σ_i. For any $(r,\tau) \in Glu_{T_0}$ there is a neighborhood U of $(\Sigma_{(r)}, u_{(r)})$ in $\mathcal{B}_A(M^+, C; g, m+\nu, (\mathbf{k}, \mathfrak{e}))$ such that $\mathcal{E}$ is trivialized over U. We choose $T_0 > R_0$ so large that for $r > T_0$

$$K_b \subset \mathcal{E}|_U.$$

Note that $u_{(r)}|_{|s_i| \leq R_0} = u_i|_{|s_i| \leq R_0}$ and *supp* $(K_b) \subset \{|s_i| \leq R_0\}$. Then $P(u_{(r)}, v)|_{|s_i| \leq R_0}$ is independent of r. Hence κ_u is well defined for any $u \in B(u_{(r)}, \epsilon)$. We can naturally identify K_b and $K_{b_{(r)}}$. We define a thickned Fredholm system and regularization equation as in **(1)**. Then there exists a neighborhood U of b, such that $D\mathcal{S}_{b'} : K_b \times W^{1,p,\alpha} \to L^{p,\alpha}$ is surjective for any $b' = (\kappa, v) \in K_b \times U$. Then $D\mathcal{S}_{b'} : K_b \times \mathcal{W}^{1,p,\alpha} \to L^{p,\alpha}$ is also surjective.

7.2 Global regularization

Denote

$$U_b^\epsilon = \{v \in \mathcal{B} |\ v = \exp_u(h + \hat{h}_0), \quad \|h\|_{1,p,\alpha} + |h_0| < \epsilon\}.$$

As $\overline{\mathcal{M}}_A(M^+, C; g_1, m_1 + \nu, (\mathbf{k}, \mathfrak{e}))$ is compact, there exist finite points b_i, $1 \leq i \leq n$ such that the collection $\{U_{b_i}^\epsilon\}$ is an open cover of $\overline{\mathcal{M}}_A(M^+, C; g_1, m_1 + \nu, (\mathbf{k}, \mathfrak{e}))$. For any $I \subset \{1, 2, ..., n\}$, setting

$$K_I = \prod_{i \in I} K_{b_i}, \quad S_I((\kappa_i)_{i \in I}, v) = \overline{\partial}_J(v) + \sum_{i \in I} \beta_{b_i} P_{u,v}(\kappa_i) \in \mathcal{E}_v.$$

One can construct a global regularization (C_I, F_I, S_I) of the original Fredholm system $(\mathcal{B}, \mathcal{E}, \overline{\partial}_J)$. From the global regularization we can obtain that (see [7,8]):

Lemma 7.1. *There exists a finite dimensional virtual orbifold system for $(\mathcal{B}, \mathcal{E}, \overline{\partial}_J)$ which is a collection of triples*

$$\{(U_I, E_I, \sigma_I) | I \subset \{1, 2, \cdots, n\}\}$$

indexed by a partially ordered set $(I = 2^{\{1,2,\cdots,n\}}, \subset)$, where

1. *$\{U_I | I \subset \{1, 2, \cdots, n\}\}$ is a finite dimensional proper étale virtual groupoid, where $U_I = S_I^{-1}(0)$,*
2. *$\{E_I\}$ is a finite rank virtual vector bundle over $\{U_I\}$,*
3. *$\{\sigma_I\}$ is a section of the virtual vector bundle $\{E_I\}$ whose zeros $\{\sigma_I^{-1}(0)\}$ form a cover of $\overline{\mathcal{M}}_A(M^+, C; g_1, m_1 + \nu, (\mathbf{k}, \mathfrak{e}))$.*

Let $U_{I,\epsilon}$ be the set of points $((\kappa_i)_{i \in I}, v) \in U_I$ such that $\sum_{i \in I} \|\kappa_i\|_{p,\alpha} \leq \epsilon$, where $I \subset \{1, 2, ..., n\}$. We define $U_{I,\epsilon}(\mathbb{R} \times \widetilde{M})$, $U_{I,\epsilon}(M^+ \cup (\mathbb{R} \times \widetilde{M}))$ in the same way.

By the same argument of Theorem 5.16 we conclude that:

Theorem 7.2. *$\{U_{I,\epsilon}(M^+)\}$ is a compact virtual manifold.*

8 Gluing theory– analysis estimates

8.1 Gluing theory for 1-nodal case

We consider the case of gluing one node in Subsection §6.1. The general cases are similar.

Recall that we have a global regularization $\{(\mathbf{C},\mathbf{F},\mathbf{S})\}$. Consider the regular Fredholm system (C_I, F_I, S_I). Let $b=((u_1,u_2),\Sigma_1\wedge\Sigma_2;j_1,j_2)\in C_I$, where (Σ_1,j_1) and (Σ_2,j_2) are smooth Riemann surfaces of genus g_1 and g_2 joining at q and $u_1:\Sigma_1\to M^+$, $u_2:\Sigma_2\to\mathbb{R}\times\widetilde{M}$ are J-holomorphic maps such that $u_i(z)$ converge to the same kT-periodic orbit $x(kTt)\subset\mathcal{F}_j$ for some j as $z\to p$.

For any $(\kappa,h,h_0)\in KerD\mathcal{S}_b$, where $h\in W^{1,p,\alpha}(\Sigma;u^*TN)$, we define

$$\|(\kappa,h)\|_{1,p,\alpha}=\|\kappa\|_{p,\alpha}+\|h\|_{1,p,\alpha},\quad \|(\kappa,h,h_0)\|=\|(\kappa,h)\|_{1,p,\alpha}+|h_0|.$$

For any $(\kappa,h_{(r)})\in KerD\mathcal{S}_{b_{(r)}}$, we define

$$\|(\kappa,h_{(r)})\|=\|\kappa\|_{p,\alpha}+\|h_{(r)}\|_{1,p,\alpha,r}.$$

By using the exponential decay of u_i one can easily prove that $u_{(r)}$ are a family of approximate J-holomorphic map, precisely the following lemma holds.

Lemma 8.1. *For any $r>R_0$, we have*

$$\|\bar{\partial}_J(u_{(r)})\|_{p,\alpha,r}\le Ce^{-(\mathfrak{c}-\alpha)r}. \tag{114}$$

The constants C in the above estimates are independent of r.

8.2 Estimates of right inverse

Lemma 8.2. *Let $D\mathcal{S}_b:K_b\times\mathcal{W}^{1,p,\alpha}\to L^{p,\alpha}$ be a Fredholm operator defined in section §7. Suppose that $D\mathcal{S}_b|_{K_b\times W^{1,p,\alpha}}:K_b\times W^{1,p,\alpha}\to L^{p,\alpha}$ is surjective. Denote by $Q_b:L^{p,\alpha}\to K_b\times W^{1,p,\alpha}$ a right inverse of $D\mathcal{S}_b$. Then $D\mathcal{S}_{b_{(r)}}$ is surjective for r large enough. Moreover, there are a right inverses $Q_{b_{(r)}}$ such that*

$$D\mathcal{S}_{b_{(r)}}Q_{b_{(r)}}=Id \tag{115}$$

$$\|Q_{b_{(r)}}\|\le C \tag{116}$$

for some constant $C>0$ independent of r.

Proof: We first construct an approximate right inverse $Q'_{b_{(r)}}$ such that the following estimates holds

$$\|Q'_{b_{(r)}}\|\le C_1 \tag{117}$$

$$\|D\mathcal{S}_{b_{(r)}}\circ Q'_{b_{(r)}}-Id\|\le\frac{1}{2}. \tag{118}$$

Then the operator $D\mathcal{S}_{b_{(r)}}\circ Q'_{b_{(r)}}$ is invertible and a right inverse $Q_{b_{(r)}}$ of $D\mathcal{S}_{b_{(r)}}$ is given by

$$Q_{b_{(r)}}=Q'_{b_{(r)}}(D\mathcal{S}_{b_{(r)}}\circ Q'_{b_{(r)}})^{-1} \tag{119}$$

Denote $\beta_1 = \beta(3/2 - \frac{Tks_1}{lr})$. Let $\beta_2 \geq 0$ be a smooth function such that $\beta_2^2 = 1 - \beta_1^2$. Given $\eta \in L_r^{2,\alpha}$, we have a pair (η_1, η_2), where

$$\eta_1 = \beta_1 \eta, \quad \eta_2 = \beta_2 \eta.$$

Let $Q_b(\eta_1, \eta_2) = (\kappa_b, h)$. We may write h as (h_1, h_2), and define

$$h_{(r)} = h_1 \beta_1 + h_2 \beta_2. \tag{120}$$

Note that on $\{\frac{lr}{2Tk} \leq s_1 \leq \frac{3lr}{2Tk}\}$, $\kappa = 0$ and on $\{|s_i| \leq \frac{lr}{2Tk}\}$ we have $u_{(r)} = u_i$, $\kappa_{(r)} = \kappa_b$, so along $u_{(r)}$ we have $\kappa_{(r)} = \kappa_b$. Then we define

$$Q'_{b_{(r)}} \eta = (\kappa_{(r)}, h_{(r)}) = (\kappa_b, h_{(r)}). \tag{121}$$

Since $|\beta_1| \leq 1$ and $|\frac{\partial \beta_1}{\partial s_1}| \leq \frac{CTk}{lr}$, (117) follows from $\|Q_b\| \leq C$. We prove (118). Since $\kappa_b + D_u h = \eta$ we have

$$D\mathcal{S}_{b_{(r)}} \circ Q'_{b_{(r)}} \eta = \eta, \qquad |s_i| \leq \frac{lr}{2Tk}. \tag{122}$$

It suffices to estimate the left hand side in the left annulus $\frac{lr}{2Tk} \leq |s_i| \leq \frac{3lr}{2Tk}$. Note that in this annulus

$$\beta_1^2 + \beta_2^2 = 1, \;\; \kappa_b = 0, \quad D_{u_i} h_i = \eta_i,$$
$$\beta_1 D_{u_1} h_1 + \beta_2 D_{u_2} h_2 = (\beta_1^2 + \beta_2^2)\eta.$$

Since near $u_1(p) = u_2(p)$ (or near the periodic orbit $x(kTt)$), $D_{u_i} = \bar{\partial}_{J_0} + S_{u_i}$, we have

$$\begin{aligned} & D\mathcal{S}_{b_{(r)}} \circ Q'_{b_{(r)}} \eta - (\beta_1^2 + \beta_2^2)\eta = \kappa_{b_{(r)}} + D_{u_{(r)}} h_{(r)} - (\beta_1^2 + \beta_2^2)\eta \\ = \;& (\bar{\partial}\beta_1) h_1 + \beta_1 (S_{u_{(r)}} - S_{u_1}) h_1 + (\bar{\partial}\beta_2) h_2 + \beta_2 (S_{u_{(r)}} - S_{u_2}) h_2. \end{aligned} \tag{123}$$

By the exponential decay of S and $\beta_1^2 + \beta_2^2 = 1$ we get

$$\begin{aligned} & \left\| D\mathcal{S}_{b_{(r)}} \circ Q'_{b_{(r)}} \eta - \eta \right\|_{p,\alpha,r} = \left\| D\mathcal{S}_{b_{(r)}} \circ Q'_{b_{(r)}} \eta - (\beta_1^2 + \beta_2^2)\eta \right\|_{p,\alpha,r} \\ & \leq \frac{C_1}{r} \left(\|h_1\|_{p,\alpha} + \|h_2\|_{p,\alpha} \right) \leq \frac{C_2}{r} \|\eta\|_{p,\alpha,r} \end{aligned} \tag{124}$$

In the last inequality we used that $\|Q_b\| \leq C$ and $(h_1, h_2) = \pi_2 \circ Q_b(\eta_1, \eta_2)$. Then (118) follows by choosing r big enough. The estimate (118) implies that

$$\frac{1}{2} \leq \|D\mathcal{S}_{b_{(r)}} \circ Q'_{b_{(r)}}\| \leq \frac{3}{2}. \tag{125}$$

Then (116) follows. □

8.3 Isomorphism between $KerD\mathcal{S}_b$ and $KerD\mathcal{S}_{b_{(r)}}$

Put

$$E_1 := KerD\mathcal{S}_{u_1}, \;\; E_2 := KerD\mathcal{S}_{u_2}, \quad \mathbb{H} = T_q(\mathcal{F}_j) \oplus (span\{\frac{\partial}{\partial a}\}),$$

Denote

$$KerD\mathcal{S}_b := E_1 \bigoplus_{\mathbb{H}} E_2.$$

For a fixed gluing parameter $(r) = (r, \tau)$ we define a map $I_r : KerD\mathcal{S}_b \longrightarrow KerD\mathcal{S}_{b_{(r)}}$. For any $(\kappa, h, h_0) \in KerD\mathcal{S}_b$, where $h \in W^{1,p,\alpha}(\Sigma; u^*TN)$, we write $h = (h_1, h_2)$, and define

$$h_{(r)} = \hat{h}_0 + h_1\beta_1 + h_2\beta_2, \tag{126}$$

$$I_r(\kappa, h, h_0) = (\kappa, h_{(r)}) - Q_{b_{(r)}} \circ D\mathcal{S}_{b_{(r)}}(\kappa, h_{(r)}). \tag{127}$$

Lemma 8.3. *I_r is an isomorphisms for r big enough.*

Proof: The proof is basically a similar gluing argument as in [9]. The proof is devides into 2 steps.

Step 1. We define a map $I'_r : KerD\mathcal{S}_{b_{(r)}} \longrightarrow KerD\mathcal{S}_b$ and show that I'_r is injective for r big enough. For any $(\kappa, h) \in KerD\mathcal{S}_{b_{(r)}}$ we denote by h_i the restriction of h to the part $|s_i| \leq \frac{lr}{kT} + \frac{1}{\alpha}$, we get a pair (h_1, h_2). Let

$$h_0 = \int_{S^1} h\left(\frac{lr}{kT}, t\right) dt. \tag{128}$$

We denote

$$\beta[h] = \left((h_1 - \hat{h}_0)\beta\left(\frac{\alpha lr}{kT} + 1 - \alpha s_1\right) + \hat{h}_0,\ (h_2 - \hat{h}_0)\beta\left(\frac{\alpha lr}{kT} + 1 + \alpha s_2\right) + \hat{h}_0\right)$$

and define $I'_r : KerD\mathcal{S}_{b_{(r)}} \longrightarrow KerD\mathcal{S}_b$ by

$$I'_r(\kappa, h) = (\kappa, \beta[h]) - Q_b \circ D\mathcal{S}_b(\kappa, \beta[h]), \tag{129}$$

where Q_b denotes the right inverse of $D\mathcal{S}_b|_{K_b \times W^{1,p,\alpha}} : K_b \times W^{1,p,\alpha} \to L^{p,\alpha}$. Since $D\mathcal{S}_b \circ Q_b = D\mathcal{S}_b|_{K_b \times W^{1,p,\alpha}} \circ Q_b = I$, we have $I'_r(KerD\mathcal{S}_{b_{(r)}}) \subset KerD\mathcal{S}_b$.

Since κ and $D_u(\beta(h - \hat{h}_0))$ have compact support and $S_u \in L^{p,\alpha}$, we have $D\mathcal{S}_b(\kappa, \beta[h]) \in L^{p,\alpha}$. Then $Q_b \circ D\mathcal{S}_b(\kappa, \beta[h]) \in K_b \times W^{1,p,\alpha}$.

Let $(\kappa, h) \in KerD\mathcal{S}_{b_{(r)}}$ such that $I'_r(\kappa, h) = 0$. Since $\beta(h - \hat{h}_0) \in W^{1,p,\alpha}$ and $Q_b \circ D\mathcal{S}_b(\kappa, \beta[h]) \in K_b \times W^{1,p,\alpha}$, then $I'_r(\kappa, h) = 0$ implies that $h_0 = 0$. From (129) we have

$$\|I'_r(\kappa, h) - (\kappa, \beta h)\|_{1,p,\alpha} \leq C_1 \|\kappa + D_u(\beta h)\|_{p,\alpha}$$

$$= C_1 \left\|\kappa + \beta\left(D_u h + D_{u_{(r)}} h + \kappa - D_{u_{(r)}} h - \kappa\right) + (\bar{\partial}\beta)h\right\|_{p,\alpha}$$

Since $(\kappa, h) \in KerD\mathcal{S}_{b_{(r)}}$, we have $\kappa + D_{u_{(r)}} h = 0$. We choose $\frac{lr}{2kT} > R_0$. As $\kappa|_{|s_i| \geq R_0} = 0$ and $\beta|_{|s_i| \leq \frac{lr}{kT}} = 1$ we have $\kappa = \beta\kappa$. Therefore

$$\left\|I'_r(\kappa, h) - (\kappa, \beta h)\right\|_{1,p,\alpha} \leq C_1 \|(\bar{\partial}\beta)h + \beta(S_u - S_{u(r)})h\|_{p,\alpha}.$$

Note that

$$S_u = S_{u(r)} \quad if \ \ s_1 \leq \frac{lr}{2kT}, \ or \ \ s_2 \geq -\frac{lr}{2kT}.$$

By exponential decay of S we have

$$\|(S_u - S_{u(r)})\beta h\|_{p,\alpha} \leq Ce^{-\mathfrak{c}\frac{lr}{2kT}} \|\beta h\|_{1,p,\alpha}$$

for some constant $C > 0$. Since $(\bar{\partial}\beta(\frac{\alpha lr}{kT}+1-\alpha s_1))h_1$ supports in $\frac{lr}{kT} \leq s_1 \leq \frac{lr}{kT}+\frac{1}{\alpha}$, and over this part

$$|\bar{\partial}\beta(\frac{\alpha lr}{kT}+1-\alpha s_1)| \leq 2|\alpha|$$

$$\beta(\frac{\alpha lr}{kT}+1+\alpha s_2)=1, \quad e^{2\alpha|s_1|} \leq e^4 e^{2\alpha|s_2|},$$

we obtain

$$\|(\bar{\partial}\beta(\frac{\alpha lr}{kT}+1-\alpha s_1))h_1\|_{p,\alpha} \leq 2|\alpha|e^4\|h_2\|_{p,\alpha} \leq 2|\alpha|e^4\|\beta h\|_{p,\alpha}.$$

Similar inequality for $(\bar{\partial}\beta(\frac{\alpha lr}{kT}+1+\alpha s_2))h_2$ also holds. So we have

$$\|(\bar{\partial}\beta)h\|_{p,\alpha} \leq 4|\alpha|e^4\|\beta h\|_{1,p,\alpha}.$$

Hence

$$\|I'_r(\kappa,h)-(\kappa,\beta h)\|_{1,p,\alpha} \leq C_3(|\alpha|+e^{-\frac{\mathfrak{c}lr}{2kT}})\|\beta h\|_{1,p,\alpha} \leq 1/2\|\beta h\|_{1,p,\alpha} \tag{130}$$

for some constant $C_3 > 0$, here we choosed $0 < \alpha < \frac{1}{4C_4}$ and r big enough such that $\frac{lr}{kT} > \frac{1}{\alpha}$ and $C_3 e^{-\frac{\mathfrak{c}lr}{2kT}} < 1/4$.

Then $I'_r(\kappa,h)=0$ and (130) gives us

$$\|\kappa\|_{p,\alpha}=0, \;\; \|\beta h\|_{1,p,\alpha}=0.$$

It follows that $\kappa=0, \;\; h=0$. So I'_r is injective.

Step 2. Since $\|Q_{b_{(r)}}\|$ is uniformly bounded, from (127) and (116), we have

$$\|I_r((\kappa,h),h_0)-(\kappa,h_{(r)})\|_{1,p,\alpha,r} \leq C\|D\mathcal{S}_{b_{(r)}}(\kappa,h_{(r)})\|.$$

By a similar culculation as in the proof of Lemma 8.2 we obtain

$$\|I_r((\kappa,h),h_0)-(\kappa,h_{(r)})\|_{1,p,\alpha,r} \leq \frac{C}{r}(\|h\|_{p,\alpha}+|h_0|). \tag{131}$$

In particular, it holds for $p=2$. It remains to show that $\|h_{(r)}\|_{2,\alpha,r}$ is close to $\|h\|_{2,\alpha}$. Denote π the projection into the second component, that is, $\pi((\kappa,h),h_0)=h$. Then $\pi(kerD\mathcal{S}_b)$ is a finite dimentional space. Let $f_i, \; i=1,..,d$ be an orthonormal basis. Then $F=\sum f_i^2 e^{2\alpha|s|}$ is an integrable function on Σ. For any $\epsilon > 0$, we may choose R_0 so big that

$$\int_{|s_i|\geq R_0} F \leq \epsilon.$$

Then the restriction of h to $|s_i| \geq R_0$ satisfies

$$\|h|_{|s_i|\geq R_0}\|_{2,\alpha} \leq \epsilon\|h\|_{2,\alpha},$$

therefore

$$\|h_{(r)}\|_{2,\alpha,r} \geq \|h|_{|s_i|\leq R_0}\|_{2,\alpha}+|h_0| \geq (1-\epsilon)\|h\|_{2,\alpha}+|h_0|, \tag{132}$$

for $r > R_0$. Suppose that $I_r((\kappa,h),h_0)=0$. Then (131) and (132) give us $h=0$ and $h_0=0$, and so $\kappa=0$. Hence I_r is injective.

The **step 1** and **step 2** together show that both I_r and I'_r are isomorphisms for r big enough. □

8.4 Gluing maps

There is a neighborhood U of $b_{(r)}$ in $\mathcal{B}_A(M^+, C; g, m+\nu, (\mathbf{k}, \mathfrak{c}))$ such that $\mathcal{E}$ is trivialized over U, more precisely, $P_{u_{(r)},v}$ gives a isomorphism $\mathcal{E}|_{u_{(r)}} \to \mathcal{E}|_v$ for any $v \in U$.

Consider a map

$$\begin{aligned} &F_{(r)} : K_{b_{(r)}} \times \mathcal{W}_r^{1,p,\alpha}(\Sigma_{(r)}; u_{(r)}^* TM^+) \to L_r^{p,\alpha}(u_{(r)}^* TM^+ \otimes \wedge^{0,1}) \\ &F_{(r)}(\kappa, h) = P_{exp_{u_{(r)}} h, u_{(r)}} \left(\bar{\partial}_J exp_{u_{(r)}} h + \kappa \right). \end{aligned}$$

Let $(\kappa_\tau, h_\tau) : [0, \delta) \longrightarrow K_{b_{(r)}} \times \mathcal{W}_r^{1,p,\alpha}(\Sigma_{(r)}; u_{(r)}^* TM^+)$ be a curve satisfying

$$\kappa_0 = \kappa, \quad h_0 = h, \quad \frac{d}{d\tau}\kappa_\tau \mid_{\tau=0} = \eta, \quad \frac{d}{d\tau} h_\tau \mid_{\tau=0} = g$$

By the same method of [23] we can prove

Lemma 8.4. 1. $dF_{(r)}(0) = D\mathcal{S}_{b_{(r)}}$.

2. *There is a constant* $\hbar_2 > 0$ *such that for any* $(\kappa, h) \in K_{b_{(r)}} \times \mathcal{W}_r^{1,p,\alpha}(\Sigma_{(r)}; u_{(r)}^* TM^+)$ *with* $\|(\kappa, h)\| < \hbar_2$ *and* $\|F_{u_{(r)}}(\kappa, h)\|_{p,\alpha,r} \leq 2\hbar_2$, *the following inequality holds*

$$\|dF_{(r)}(\kappa, h) - dF_{(r)}(0)\| \leq \frac{1}{4\mathcal{C}_2}. \tag{133}$$

Let $\hbar_3 = \min\{\hbar_1, \hbar_2\}$. Note that $M^+ = M_0^+ \times ([0, \infty) \times \widetilde{M})$. By the definition of the metrics $\langle , \rangle$ (see (16) and (17)) the curvature $|Rm|$ is uniform bounded above. Then there exists a constant $C_2 > 0$ such that $|dexp_p| \leq C_2$ for any $p \in M^+$. For any $\|(\kappa, h)\| \leq \frac{\hbar_3}{8C_2\mathcal{C}_2}$,

$$\begin{aligned} \|F_{(r)}(\kappa, h)\|_{p,\alpha,r} &= \|\bar{\partial}_J exp_{u_{(r)}} h + \kappa\|_{p,\alpha,r} \\ &\leq \|\overline{\partial}_J u_{(r)}\|_{p,\alpha,r} + \|dexp_{u_{(r)}} \partial_s h\|_{p,\alpha,r} + \|dexp_{u_{(r)}} \partial_t h\|_{p,\alpha,r} + \|\kappa\|_{p,\alpha} \\ &\leq \|\overline{\partial}_J u_{(r)}\|_{p,\alpha,r} + C_2\|(\kappa, h)\| \leq C_1 e^{-(\mathfrak{c}-\alpha)r} + \frac{\hbar_3}{8\mathcal{C}_2} \leq \frac{\hbar_3}{4\mathcal{C}_2} \end{aligned} \tag{134}$$

when r big enough. It follows from (2) of Lemma 8.4 and (134) that

$$\|dF_{(r)}(\kappa, h) - D\mathcal{S}_{b_{(r)}}\| \leq \frac{1}{4\mathcal{C}_2}. \tag{135}$$

Then $F_{(r)}$ satisfies the conditions in Lemma 2.10. It follows that for any $(\kappa, h) \in kerD\mathcal{S}_{b_{(r)}}$ with $\|(\kappa, h)\| \leq \frac{\hbar_3}{8C_2\mathcal{C}_2}$ there exists a unique (κ_v, ζ) such that $F_{(r)}(\kappa_v, \zeta) = 0$. Since $K_{b_{(r)}} \times \mathcal{W}_r^{1,p,\alpha}(\Sigma_{(r)}; u_{(r)}^* TM^+) = imQ_{b_{(r)}} + kerD\mathcal{S}_{b_{(r)}}$ and $Q_{b_{(r)}}$ is injective, then there exists a unique smooth map

$$f_{(r)} : Ker\ D\mathcal{S}_{b_{(r)}} \to L_r^{p,\alpha}(u_{(r)}^* TM^+ \otimes \wedge^{0,1})$$

such that $(\kappa_v, \zeta) = ((\kappa, h) + Q_{b_{(r)}} \circ f_{(r)}(\kappa, h))$.

On the other hand, suppose that (κ_v, ζ) satisfies

$$F_{(r)}(\kappa_v, \zeta) = 0, \qquad \|(\kappa_v, \zeta)\| \leq \frac{\hbar_3}{8(1 + \mathcal{C}_2\|D\mathcal{S}_{b_{(r)}}\|)C_2\mathcal{C}_2}. \tag{136}$$

We write $(\kappa_v, \zeta) = (\kappa, h) + Q_{b_{(r)}} \circ \eta$ where $(\kappa, h) \in kerD\mathcal{S}_{b_{(r)}}$ and $\eta \in L_r^{p,\alpha}(u_{(r)}^* TM^+ \otimes \wedge^{0,1})$. Since $(\kappa, h) = (I - Q_{b_{(r)}} \circ D\mathcal{S}_{b_{(r)}})(\kappa_v, \zeta)$, we have $\|(\kappa, h)\| \leq \frac{\hbar_3}{8C_2\mathcal{C}_2}$, then $\eta = f_{(r)}(\kappa, h)$, i.e.,

$$(\kappa_v, \zeta) = (\kappa, h) + Q_{b_{(r)}} \circ f_{(r)}(\kappa, h)). \tag{137}$$

Hence the zero set of $F_{(r)}$ is locally the form $((\kappa, h), Q_{b_{(r)}} \circ f_{(r)}(\kappa, h))$, i.e

$$F_{(r)}((\kappa, h) + Q_{b_{(r)}} \circ f_{(r)}(\kappa, h)) = 0 \tag{138}$$

where $(\kappa, h) \in Ker\ D\mathcal{S}_{b_{(r)}}$.

For fixed (r) denote

$$U_{b_{(r)}} = \left\{ (\kappa, h) \in K_{b_{(r)}} \times \mathcal{W}^{1,p,\alpha,r}(\Sigma_{(r)}; u_{(r)}^* TM^+) \mid \bar{\partial}_J exp_{u_{(r)}} h + \kappa = 0 \right\},$$

$$(\kappa_{(r)}, \zeta_{(r)}) := I_r(\kappa, \zeta) + Q_{b_{(r)}} \circ f_{(r)} \circ I_r(\kappa, \zeta)).$$

Define π_1, π_2 by

$$\pi_1(\kappa, h) = \kappa, \quad \text{and} \quad \pi_2(\kappa, h) = h, \quad \text{for any } (\kappa, h).$$

Since $I_r : kerD\mathcal{S}_b \to KerD\mathcal{S}_{b_{(r)}}$ is a isomorphism, we have proved the following

Lemma 8.5. *There is a neighborhood O of 0 in $kerD\mathcal{S}_b$ and a neighborhood O_j of (j_1, j_2) and $R_0 > 0$ such that*

$$glu_{(r)} : O_j \times O \times \mathbb{Z}_k \to U_{b_{(r)}}$$

defined by

$$glu_{(r)}(j_1, j_2, \kappa, \zeta, n) = (\pi_1(\kappa_{(r)}, \zeta_{(r)}), \exp_{u_{(r)}}(\pi_2(\kappa_{(r)}, \zeta_{(r)})))$$

where for $r > R_0$ is a family of orientation preserving local diffeomorphisms.

We may choose $((j_1, j_2), r, \tau, \kappa, \zeta)$ as a local coordinate system around b in $U_{I,\epsilon}$. We write $f_{(r)} \circ I_r$ as a function $f(j_1, j_2, r, \tau, \kappa, \zeta)$. As I_r is a smooth map we can see that $f(j_1, j_2, r, \tau, \kappa, \zeta)$ is smooth (see p.131 in [30]).

8.5 Surjectivity and injectivity

Proposition 8.6. *The gluing map in Lemma 8.5 is surjective and injective in the sense of the Gromov-Uhlenbeck convergence.*

Proof. The injectivity follows immediately from the implicit function theorem. We prove the surjectivity. Let $\Gamma = ((\kappa_v, v), \Sigma_{(r)}) \in U_{b_{(r)}}$ and $\zeta \in W_r^{1,p,\alpha}$ such that $\exp_{u_{(r)}}(\zeta) = v$. Suppose that $\|\kappa_v\|_{p,\alpha} + |\zeta|_{C^1(\mathfrak{D}_i(R_0))} \leq \epsilon$. By the same argument of Lemma 6.1 we have (κ, ζ) satisfies (136). Then there exists a unique $(\kappa, h) \in Ker\ D\mathcal{S}_b$ such that

$$(\kappa_v, \zeta) = I_r(\kappa, h) + Q_{b_{(r)}} \circ f_{(r)} \circ I_r(\kappa, h)).$$

The surjectivity follows.

9 Estimates of differentiations for gluing parameters

In the section, let $\alpha << \mathfrak{c}$ be a constant. We estimate the differentiations for gluing parameters. This is another key point of this paper.

9.1 Linear analysis on weight sobolov spaces

Let $\Sigma = \mathbb{R} \times S^1$, $E = \Sigma \times \mathbb{R}^{2n}$. Let $\pi : E \longrightarrow \Sigma$ be the trivial vector bundle. Consider the Cauchy-Riemannian operator D on E defined by

$$D = \frac{\partial}{\partial s} + J_0 \frac{\partial}{\partial t} + S = \bar{\partial}_{J_0} + S,$$

where $S(s,t) = (S_{ij}(s,t))_{2n\times 2n}$ is a matrix such that for any $k > 0$,

$$\sum_{i+j=k} \left| \frac{\partial^k S}{\partial^i s \partial^j t} \right| \leq C_k e^{-\mathfrak{c}s}. \tag{139}$$

for some constant $C_k > 0$. Therefore, the operator $H_s = J_0 \frac{d}{dt} + S$ converges to $H_\infty = J_0 \frac{d}{dt}$ as $s \to \infty$.

We choose a small weight α for each end such that $H_\infty - \alpha = J_0 \frac{d}{dt} - \alpha$ is invertible. Let W be the function in section §3. We can define the space $W^{1,p,\alpha}(\Sigma; E)$ and $L^{p,\alpha}(E \otimes \wedge^{0,1})$. Then the operator $D : W^{1,p,\alpha}(\Sigma; E) \to L^{p,\alpha}(E \otimes \wedge^{0,1})$ is a Fredholm operator so long as α does not lie in the spectrum of the operator $H_{i\infty}$ for all $i = 1, \cdots, \nu$.

By the observation of Donaldson we know that the multiplication by W gives isometries from $W^{1,2,\alpha}$ to $W^{1,2}$ and $L^{2,\alpha}$ to L^2, and

$$\|f\|_{2,\alpha} = \|Wf\|_{L^2}, \quad C^{-1}\|Wf\|_{W^{1,2}} \leq \|f\|_{1,2,\alpha} \leq C\|Wf\|_{W^{1,2}}. \tag{140}$$

Consider the operator

$$D : W^{1,2,\alpha} \to L^{2,\alpha}.$$

It is equivalent to the map (see [9],P59)

$$D_W := D - \frac{\partial}{\partial s} \log W = WDW^{-1} : W^{1,2} \to L^2.$$

Let $h \in W^{1,2,\alpha}$ be the solution of the equation $Dh = \eta$, where $\eta \in L^{2,\alpha}$. Obviously,

$$D_W(Wh) = WD(W^{-1}Wh) = W\eta.$$

Let $\rho = W\eta$, $f = Wh$. Then f satisfies the equation $D_W f = \rho$.

When $s \geq R_0$ we consider the operator $D - \alpha : W^{1,2} \to L^2$. We write $D := D - \alpha = \frac{\partial}{\partial s} + L + S$, where $L := J_0 \frac{\partial}{\partial t} - \alpha$ is a invertible elliptic operator on S^1 when $0 < \alpha < 1$.

The space $L^2(S^1)$ can be decomposed two infinite spaces H^+ and H^-(see P33 in [9]), where

$$H^+ = \left\{ \sum a_n e^{int} | n \geq 0 \right\}, \quad H^- = \left\{ \sum a_n e^{int} | n < 0 \right\}.$$

Since L is an invertible operator, there exists a complete eigen-function space decomposition such that $L\phi_\lambda = \lambda\phi_\lambda$ where $|\lambda| > \delta$ for some $\delta > 0$. Obviously, $\lambda \in \mathbb{Z} - \alpha$. By choosing $0 < \alpha < 1/2$ we have $\alpha = \min |\lambda|$.

Using the same method of Donaldson in [9] we can prove the following lemmas (for details please see our next paper [20].)

For $A > 1$ we consider the finite tube $(-A, A) \times S^1$ and denote

$$B_A^- = (-A, -A+1) \times S^1, \quad B_A^+ = (A-1, A) \times S^1.$$

Lemma 9.1. *Let h be a solution of $Dh = 0$ over the finite tube $(-A, A) \times S^1$. Then for any $0 < \alpha < \min\{\mathfrak{c}, \frac{1}{2}\}$, there exists a constant $C_1 > 0$ such that for any $0 < |s_o| + 1 < A$*

$$\|h \mid_{(-|s_o|+1, |s_o|-1) \times S^1} \|_{1,p,\alpha} \leq C_1 e^{-\alpha(A-|s_o|)} \left(\|h|_{B_A^+}\|_{2,\alpha} + \|h|_{B_A^-}\|_{2,\alpha} \right). \tag{141}$$

Using this lemma we can prove the following corollary.

Corollary 9.2. *Suppose that D is surjective. Denote by Q a bounded right inverse of D. Let $h = Q\eta$ be a solution of $Dh = \eta$ over the finite tube $(0, \frac{2lr}{kT}) \times S^1$. Then for any $0 < \alpha < \min\{\mathfrak{c}, \frac{1}{2}\}$, there exists a constant $C_2 > 0$ such that for any $0 < A < \frac{lr}{2kT}$,*

$$\|h \mid_{(\frac{lr}{2kT}, \frac{3lr}{2kT}) \times \mathbb{S}^1} \|_{1,2,\alpha} \leq C_2 \left(e^{-\alpha A} \|\eta\|_{2,\alpha} + \|\eta \mid_{\frac{lr}{2kT} - A \leq s \leq \frac{3lr}{2kT} + A} \|_{2,\alpha} \right). \tag{142}$$

Proof. Denote by $\hat{\eta}$ the restriction of η to the part $\frac{lr}{2kT} - A \leq s \leq \frac{3lr}{2kT} + A$. Let $\hat{h} = Q\hat{\eta}$. Then $D(h - \hat{h}) = 0$ in $\frac{lr}{2kT} - A \leq s \leq \frac{3lr}{2kT} + A$. By (141) and

$$\|\hat{\eta}\|_{1,p,\alpha} = \|\eta \mid_{\frac{lr}{2kT} - A \leq s \leq \frac{3lr}{2kT} + A} \|_{2,\alpha} \qquad \|\hat{h}\|_{1,p,\alpha} = \|Q\hat{\eta}\|_{1,p,\alpha} \leq C \|\hat{\eta}\|_{1,p,\alpha}$$

we have

$$\begin{aligned} \|h \mid_{(\frac{lr}{2kT}, \frac{3lr}{2kT}) \times \mathbb{S}^1} \|_{1,2,\alpha} &\leq \|h - \hat{h} \mid_{(\frac{lr}{2kT}, \frac{3lr}{2kT}) \times \mathbb{S}^1} \|_{1,2,\alpha} + \|\hat{h} \mid_{(\frac{lr}{2kT}, \frac{3lr}{2kT}) \times \mathbb{S}^1} \|_{1,2,\alpha} \\ &\leq C_1 \left(e^{-\alpha A} \|h - \hat{h} \mid_{\frac{lr}{2kT} - A \leq s \leq \frac{3lr}{2kT} + A} \|_{1,2,\alpha} + \|\eta \mid_{\frac{lr}{2kT} - A \leq s \leq \frac{3lr}{2kT} + A} \|_{2,\alpha} \right). \end{aligned}$$

Then the corollary follows from $\|h - \hat{h}\|_{1,2,\alpha} = \|Q(\eta - \hat{\eta})\|_{1,2,\alpha} \leq C\|\eta\|_{2,\alpha}$. $\Box$

9.2 Estimates of $\frac{\partial Q_{b_{(r)}}}{\partial r}$

Lemma 9.3. *Let $D\mathcal{S}_b : K_b \times \mathcal{W}^{1,p,\alpha} \to L^{p,\alpha}$ be a Fredholm operator. Suppose that $D\mathcal{S}_b|_{K_b \times W^{1,p,\alpha}} : K_b \times W^{1,p,\alpha} \to L^{p,\alpha}$ is surjective. Denote by $Q_b : L^{p,\alpha} \to K_b \times W^{1,p,\alpha}$ a right inverse of $D\mathcal{S}_b$. Let $Q_{b_{(r)}}$ be an right inverse of $D\mathcal{S}_{b_{(r)}}$ defined in (119). Then there exists a constant $\mathcal{C}_3 > 0$, independent of r, such that $\left\| \frac{\partial}{\partial r} Q_{b_{(r)}} \right\| \leq \mathcal{C}_3$ and for any $0 < s_0 \leq \frac{lr}{4kT}$*

$$\left\| \frac{\partial}{\partial r} \left(Q_{b_{(r)}} \eta_r \right) \right\| \leq \mathcal{C}_3 \left[\| \frac{\partial}{\partial r} \eta_r \|_{p,\alpha,r} + \frac{1}{r} \|\eta_r \mid_{\frac{lr}{4kT} \leq |s_i| \leq \frac{7lr}{4kT}} \|_{p,\alpha,r} + e^{-\alpha \frac{lr}{8kT}} \|\eta_r\|_{2,\alpha,r} \right], \tag{143}$$

for any $\eta_r \in L_r^{2,\alpha}$.

Proof. Given $\eta_r \in L_r^{2,\alpha}$, denote $\tilde{\eta}_r = (D\mathcal{S}_{b_{(r)}} \circ Q'_{b_{(r)}})^{-1} \eta_r$, we have a pair $(\tilde{\eta}_1, \tilde{\eta}_2)$, where $\tilde{\eta}_1 = \beta_1 \tilde{\eta}_r$, $\tilde{\eta}_2 = \beta_2 \tilde{\eta}_r$. Set

$$Q_b(\tilde{\eta}_1, \tilde{\eta}_2) = (\tilde{\kappa}_b, \tilde{h}_1, \tilde{h}_2), \;\; \tilde{h}_{(r)} = \tilde{h}_1 \beta_1 + \tilde{h}_2 \beta_2.$$

Then $Q_{b_{(r)}}\eta_r = Q'_{b_{(r)}}\tilde{\eta}_r = (\tilde{\kappa}_b, \tilde{h}_{(r)})$. So

$$\frac{\partial}{\partial r}\left(Q_{b_{(r)}}\eta_r\right)|_{|s_i|\leq\frac{lr}{2kT}} = \left(\frac{\partial\tilde{\kappa}_b}{\partial r}, \sum\frac{\partial\tilde{h}_i}{\partial r}\beta_i + \sum\tilde{h}_i\frac{\partial\beta_i}{\partial r}\right)|_{|s_i|\leq\frac{lr}{2kT}} = Q_b\left(\frac{\partial}{\partial r}\tilde{\eta}_1, \frac{\partial}{\partial r}\tilde{\eta}_2\right)|_{|s_i|\leq\frac{lr}{2kT}},$$

where we used $\beta\,|_{|s_i|\leq\frac{lr}{2kT}} = 1$. As $\frac{\partial}{\partial r}\tilde{\eta}_i = \beta_i\frac{\partial}{\partial r}\tilde{\eta}_r + \frac{\partial\beta_i}{\partial r}\tilde{\eta}_r$, we have

$$\left\|\left(\frac{\partial\kappa}{\partial r}, \frac{\partial\tilde{h}_1}{\partial r}, \frac{\partial\tilde{h}_2}{\partial r}\right)\right\| = \left\|Q_b\left(\frac{\partial}{\partial r}\tilde{\eta}_1, \frac{\partial}{\partial r}\tilde{\eta}_2\right)\right\| \leq C_1\left\|\frac{\partial\tilde{\eta}_r}{\partial r}\right\|_{p,\alpha,r} + \frac{C_1}{r}\left\|\tilde{\eta}_r\,|_{\frac{lr}{2kT}\leq|s_i|\leq\frac{3lr}{2kT}}\right\|_{p,\alpha,r}. \tag{144}$$

By the same calculation as (123) we have

$$\begin{aligned} \eta_r - \tilde{\eta}_r &= D\mathcal{S}_{b_{(r)}}\circ Q'_{b_{(r)}}\tilde{\eta}_r - (\beta_1^2+\beta_2^2)\tilde{\eta}_r = \kappa_{b_{(r)}} + D_{u_{(r)}}\tilde{h}_{(r)} - (\beta_1^2+\beta_2^2)\tilde{\eta}_r \\ &= \sum(\bar{\partial}\beta_i)\tilde{h}_i + \sum\beta_i(S_{u_{(r)}} - S_{u_i})\tilde{h}_i. \end{aligned} \tag{145}$$

By the exponential decay $S_{u_{(r)}}, S_{u_i}$ and

$$\bar{\partial}\beta|_{|s_i|\leq\frac{lr}{2kT}} = 0,\ |\bar{\partial}\beta| \leq \frac{C}{r}, S_{u_{(r)}}|_{|s_i|\leq\frac{lr}{2kT}} = S_{u_i}|_{|s_i|\leq\frac{lr}{2kT}}, \tag{146}$$

we conclude from (145) that

$$\|\tilde{\eta}_r\,|_{\frac{lr}{4kT}\leq|s_i|\leq\frac{7lr}{4kT}}\|_{2,\alpha,r} \leq \left\|\eta_r\,|_{\frac{lr}{4kT}\leq|s_i|\leq\frac{7lr}{4kT}}\right\|_{2,\alpha,r} + \frac{C_2}{r}\|\tilde{h}_i\,|_{\frac{lr}{2kT}\leq|s_i|\leq\frac{3lr}{2kT}}\|_{1,2,\alpha,r}. \tag{147}$$

Applying Corollary 9.2 to (147) we have

$$\left(1-\frac{C_3}{r}\right)\|\tilde{\eta}_r\,|_{\frac{lr}{4kT}\leq|s_i|\leq\frac{7lr}{4kT}}\|_{2,\alpha,r} \leq \left\|\eta_r\,|_{\frac{lr}{4kT}\leq|s_i|\leq\frac{7lr}{4kT}}\right\|_{2,\alpha,r} + e^{-\alpha\frac{lr}{4kT}}\|\eta_r\|_{2,\alpha,r}, \tag{148}$$

where we used $\|\tilde{\eta}_r\|_{p,\alpha,r} \leq 2\|\eta_r\|_{p,\alpha,r}$. Taking the derivative of (145), by the exponential decay of $S_{u_{(r)}}, S_{u_i}$, and $|\frac{\partial}{\partial r}\bar{\partial}\beta| \leq \frac{C}{r^2}$, using (146), (142) and (145), we have

$$\begin{aligned} \|\frac{\partial}{\partial r}\tilde{\eta}_r\|_{p,\alpha,r} &\leq \|\frac{\partial}{\partial r}\eta_r\|_{p,\alpha,r} + \frac{C_4}{r^2}\|\tilde{h}_i\,|_{\frac{lr}{2kT}\leq|s_i|\leq\frac{3lr}{2kT}}\|_{1,p,\alpha} + \frac{C_4}{r}\left\|\frac{\partial\tilde{h}_i}{\partial r}\,|_{\frac{lr}{2kT}\leq|s_i|\leq\frac{3lr}{2kT}}\right\|_{1,p,\alpha} \\ &\leq \|\frac{\partial}{\partial r}\eta_r\|_{p,\alpha,r} + \frac{C_5}{r^2}\|\tilde{\eta}_r\,|_{\frac{lr}{4kT}\leq|s_i|\leq\frac{7lr}{4kT}}\|_{1,p,\alpha} + e^{-\alpha\frac{lr}{4kT}}\|\eta_r\|_{2,\alpha} + \frac{C_5}{r}\left\|\frac{\partial\tilde{\eta}_r}{\partial r}\right\|_{p,\alpha,r}. \end{aligned} \tag{149}$$

Following from (149) and (148) we obtain

$$\begin{aligned} &C_1\|\frac{\partial}{\partial r}\tilde{\eta}_r\|_{p,\alpha,r} + \frac{C_1}{r}\|\tilde{\eta}_r\,|_{\frac{lr}{4kT}\leq|s_i|\leq\frac{7lr}{4kT}}\|_{2,\alpha,r} \\ &\leq C_6\left[\|\frac{\partial}{\partial r}\eta_r\|_{p,\alpha,r} + \frac{1}{r}\left\|\eta_r\,|_{\frac{lr}{4kT}\leq|s_i|\leq\frac{7lr}{4kT}}\right\|_{2,\alpha,r} + e^{-\alpha\frac{lr}{4kT}}\|\eta_r\|_{2,\alpha}\right]. \end{aligned} \tag{150}$$

(144) and (150) gives us

$$\left\|\frac{\partial}{\partial r}\left(Q_{b_{(r)}}\eta_r\right)|_{|s_i|\leq s_0}\right\|_{1,2,\alpha,r} \leq \|\frac{\partial}{\partial r}\eta_r\|_{p,\alpha,r} + \frac{C_7}{r}\|\eta_r\,|_{\frac{lr}{4kT}\leq|s_i|\leq\frac{7lr}{4kT}}\|_{1,p,\alpha} + e^{-\alpha\frac{lr}{4kT}}\|\eta_r\|_{2,\alpha}. \tag{151}$$

Using (142) and (148), we have

$$\begin{aligned}\|\sum \tilde{h}_i \tfrac{\partial \beta_i}{\partial r}\|_{1,p,\alpha} &\leq \frac{C_8}{r}\left(e^{-\alpha \frac{lr}{8kT}}\|\tilde{\eta}\|_{1,2,\alpha} + \|\tilde{\eta}\mid_{\frac{3lr}{8kT}\leq |s_i| \leq \frac{13lr}{8kT}}\|_{2,\alpha}\right)\\ &\leq \frac{C_9}{r}\left(e^{-\alpha \frac{lr}{8kT}}\|\eta\|_{1,2,\alpha} + \|\eta\mid_{\frac{lr}{4kT}\leq |s_i| \leq \frac{7lr}{4kT}}\|_{2,\alpha}\right) \end{aligned} \tag{152}$$

Note that $\frac{\partial}{\partial r}\left(Q_{b_{(r)}}\eta_r\right) = \left(\frac{\partial \tilde{\kappa}_b}{\partial r}, \sum \frac{\partial \tilde{h}_i}{\partial r}\beta_i + \sum \tilde{h}_i \frac{\partial \beta_i}{\partial r}\right)$. Then (143) follows from (144), (150) and (152). □

9.3 Estimates of $\frac{\partial I_{(r)}}{\partial r}$

Lemma 9.4. *Let $I_r : KerD\mathcal{S}_b \longrightarrow KerD\mathcal{S}_{b_{(r)}}$ be an isomorphisms defined in (127). Then there exists a constant $C > 0$, independent of r, such that*

$$\left\|\frac{\partial}{\partial r}\left(I_r(\kappa, h, h_0)\right)\right\|_{1,2,\alpha,r} \leq \frac{C}{r}\|(\kappa, h)\mid_{\frac{lr}{2kT}\leq |s_i| \leq \frac{3lr}{2kT}}\|_{1,2,\alpha} \leq \frac{C}{r}e^{-(\mathfrak{c}-\alpha)\frac{lr}{2kT}}\|(\kappa, h)\|, \tag{153}$$

for any $(\kappa, h, h_0) \in KerD\mathcal{S}_b$.

Proof. By definition κ is independent of r. Then $\frac{\partial}{\partial r}h_{(r)} = \sum \frac{\partial}{\partial r}\beta_i h_i$ and

$$\frac{\partial}{\partial r}I_r(\kappa, h, h_0) = (0, \frac{\partial h_{(r)}}{\partial r}) + \frac{\partial}{\partial r}\left(Q_{b_{(r)}}\right) D\mathcal{S}_{b_{(r)}}(\kappa, h_{(r)}) + Q_{b_{(r)}}\frac{\partial}{\partial r}\left(D\mathcal{S}_{b_{(r)}}(\kappa, h_{(r)})\right).$$

A direct calculation gives us

$$D\mathcal{S}_{b_{(r)}}(\kappa, h_{(r)}) = (\bar{\partial}\beta_1)h_1 + \beta_1(S_{u_{(r)}} - S_{u_1})h_1 + (\bar{\partial}\beta_2)h_2 + \beta_2(S_{u_{(r)}} - S_{u_2})h_2. \tag{154}$$

By (146) we have *supp* $D\mathcal{S}_{b_{(r)}}(\kappa, h_{(r)}) \subset \{\frac{lr}{2kT} \leq |s_i| \leq \frac{3lr}{2kT}\}$. Since $\left(|S_{u_{(r)}}| + |S_u|\right)\mid_{\frac{lr}{2kT}\leq |s_i| \leq \frac{3lr}{2kT}} \leq Ce^{-\mathfrak{c}\frac{lr}{2kT}}$, the lemma follows from $\|Q_{b_{(r)}}\| \leq C$, $\|\frac{\partial}{\partial r}Q_{b_{(r)}}\| \leq C$. □

9.4 Estimates of $\frac{\partial}{\partial r}\left[I_r(\kappa, \zeta) + Q_{b(r)}f_{(r)}(I_r(\kappa, \zeta))\right]$

First we prove the following lemma.

Lemma 9.5. *Let $(\kappa, \zeta) \in kerD\mathcal{S}_b$ with $\|(\kappa, \zeta)\| \leq \hbar_3$ such that*

$$(\kappa, h_{(r)}) = I_r(\kappa, \zeta) + Q_{b_{(r)}} \circ f_r \circ I_r(\kappa, \zeta) \tag{155}$$

$$\bar{\partial}_J \exp_{u_{(r)}}(h_{(r)}) + \kappa = 0. \tag{156}$$

Then there exists a positive constants C such that

$$\|f_r \circ I_r(\kappa, \zeta)|_{\frac{lr}{4kT}\leq |s_i| \leq \frac{7lr}{4kT}}\|_{p,\alpha,r} \leq Ce^{-(\mathfrak{c}'-\alpha)\frac{lr}{4kT}}\|(\kappa, \zeta)\|, \quad \forall\ \frac{lr}{8kT} \geq R_0. \tag{157}$$

Proof. From (155) and $\kappa|_{|s_i|\geq R_0} = 0$ we have

$$f_{(r)}(I_r(\kappa, \zeta)) = \left(\frac{\partial}{\partial s} + J_0\frac{\partial}{\partial t}\right)h_{(r)} + S_{u_{(r)}}h_{(r)}, \qquad \bar{\partial}_J \exp_{u_{(r)}} h_{(r)} = 0, \tag{158}$$

for any $|s_i| \geq R_0$. By the assumption u_i converges to kT-periodic orbit $x(kTt)$ as $z \to p$. In the given local pseudo-Dauboux coordinate $(a, \vartheta, \mathbf{w})$, we denote

$$\exp_{u_{(r)}}(h_{(r)}) = (\hat{a}_{(r)}(s,t), \hat{\vartheta}_{(r)}(s,t), \hat{\mathbf{w}}_{(r)}(s,t)).$$

By Theorem 2.7, (91) and (92) we have for any $|s_i| \geq R_0$

$$|\nabla(a_{(r)} - kTs_i)| \leq Ce^{-\mathfrak{c}\frac{lr}{4kT}}, \quad |\nabla(\vartheta_{(r)} - kt_i)| \leq Ce^{-\mathfrak{c}\frac{lr}{4kT}}, \tag{159}$$

$$|\nabla(\mathbf{w}_{(r)})| \leq Ce^{-\mathfrak{c}\frac{lr}{4kT}}. \tag{160}$$

Let

$$a^{\diamond}_{(r)} = a_{(r)} - kTs_i, \;\; \vartheta^{\diamond}_{(r)} = \vartheta_{(r)} - kt_i, \quad \hat{a}^{\diamond}_{(r)} = \hat{a}_{(r)} - kTs_i, \;\; \hat{\vartheta}^{\diamond}_{(r)} = \hat{\vartheta}_{(r)} - kt_i.$$

By (159) and (160) we have

$$\int_{|s_i| \geq s_o} \|\nabla a^{\diamond}_{(r)}\|^2 + \|\nabla \vartheta^{\diamond}_{(r)}\|^2 + \|\nabla \mathbf{w}_{(r)}\|^2 dsdt \leq Ce^{-\mathfrak{c}s_o}, \qquad \forall s_o \geq R_0. \tag{161}$$

Since $\|(\kappa, h_{(r)})\| \leq C\hbar_3$ and $\kappa|_{|s_i| \geq R_0} = 0$, we have

$$\|h_{(r)}|_{|s_i| \geq R_0}\|_{1,2,\alpha} \leq C\hbar_3. \tag{162}$$

By the elliptic regularity we can get

$$|h_{(r)}| \leq Ce^{-\alpha \min\{|s_1|,|s_2|\}}, \quad \forall |s_i| \geq R_0.$$

Then $\hat{\mathbf{w}}_{(r)}(|s_i| \geq R_0)$ is also in the pseudo-Dauboux coordinates when $\frac{lr}{8kT} > R_0$ and $\hbar_3$ small. By (161)

$$\int_{|s_i| \geq s_o} \|\nabla \hat{a}^{\diamond}_{(r)}\|^2 + \|\nabla \hat{\vartheta}^{\diamond}_{(r)}\|^2 + \|\nabla \hat{\mathbf{w}}_{(r)}\|^2 dsdt \leq Ce^{-\alpha s_o}, \qquad \forall s_o \geq R_0. \tag{163}$$

Then by Theorem 2.9 we have for any $\frac{lr}{4kT} \leq |s_i| \leq \frac{7lr}{4kT}$

$$|\nabla(\hat{a}_{(r)} - kTs_i)| \leq Ce^{-\mathfrak{c}\frac{lr}{4kT}}, \quad |\nabla(\hat{\vartheta}_{(r)} - kt_i))| \leq Ce^{-\mathfrak{c}\frac{lr}{4kT}}, \tag{164}$$

$$|\nabla(\hat{\mathbf{w}}_{(r)})| \leq Ce^{-\mathfrak{c}\frac{lr}{4kT}}. \tag{165}$$

Since $\nabla \exp_{u_{(r)}}(h_{(r)}) = P_{u_{(r)}, \exp_{u_{(r)}}(h_{(r)})}(\nabla u_{(r)}) + d\exp_{u_{(r)}}(\nabla h_{(r)})$, (159), (160), (164) and (165) gives us

$$|d\exp_{u_{(r)}} \nabla h_{(r)}|_{\frac{lr}{4kT} \leq |s_i| \leq \frac{7lr}{4kT}}| \leq Ce^{-\mathfrak{c}\frac{lr}{4kT}}. \tag{166}$$

Note that $d\exp^{-1}_{u_{(r)}}$ is uniform bounded as $\|h_{(r)}\|_{1,p,\alpha,r}$ small. Then

$$\|\nabla h_{(r)}|_{\frac{lr}{4kT} \leq |s_i| \leq \frac{7lr}{4kT}}\|_{p,\alpha} \leq Ce^{-(\mathfrak{c}-\alpha)\frac{lr}{4kT}}. \tag{167}$$

The lemma follows from (158), (167) and the exponential decay of $S_{u_{(r)}}$. $\square$

Lemma 9.6. *There exists a constant $C > 0$ such that,*

$$\left\| \frac{\partial}{\partial r}\left(I_r(\kappa, \zeta) + Q_{b_{(r)}} \circ f_{(r)} \circ I_r(\kappa, \zeta) \right) \right\|_{1,2,\alpha,r} \leq Ce^{-\alpha\frac{lr}{8kT}} \|(\kappa, \zeta\|_{1,2,\alpha}. \tag{168}$$

Restricting to the compact set $\{|s_i| \leq R_0\}$, we have

$$\left| \frac{\partial}{\partial r}\left(I_r(\kappa, \zeta) + Q_{b_{(r)}} \circ f_{(r)} \circ I_r(\kappa, \zeta) \right) \right| \leq Ce^{-\alpha\frac{lr}{8kT}} \|\zeta\|_{L^\infty}. \tag{169}$$

Proof. Without loss of generality, we will fix a complex structure (j_1, j_2) to simplify notations. We write

$$I_r(\kappa,\zeta) + Q_{b(r)} \circ f_{(r)}(I_r(\kappa,\zeta)) = (\kappa_r, h_{(r)}) \tag{170}$$

where $\kappa_r \in K_{b(r)}, h_{(r)} \in C^{\infty}(\Sigma; u^{\star}_{(r)}TM^{+})$. We have

$$\bar{\partial}_J exp_{u_{(r)}} h_{(r)} + \kappa_r = 0. \tag{171}$$

To simplify the expressions of formulas we denote

$$\frac{\partial}{\partial r}\exp_{u_{(r)}}(h_{(r)}) = G, \qquad \frac{\partial}{\partial r}\kappa_r = \Psi.$$

Taking differentiation of (171) with respect to r we get

$$D_{\exp_{u_{(r)}}(h_{(r)})} G + \Psi = 0. \tag{172}$$

We may write (172) as

$$P_{exp_{u_{(r)}}h_{(r)}}\left(D\mathcal{S}_{exp_{u_{(r)}}h_{(r)}}(\Psi, G)\right) - D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}\left(\kappa_r, h_{(r)}\right)\right) \tag{173}$$

$$+ D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}\left(\kappa_r, h_{(r)}\right)\right) = 0.$$

We estimate the difference

$$P_{exp_{u_{(r)}}h_{(r)}}\left(D\mathcal{S}_{exp_{u_{(r)}}h_{(r)}}(\Psi, G)\right) - D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}\left(\kappa_r, h_{(r)}\right)\right).$$

We choose $\|\eta\|_{1,p,\alpha,r}$ very small. From the Implicity function Theorem we have

$$\|f_{(r)}(I_r(\kappa,\zeta))\|_{p,\alpha,r} \le C\|(\kappa,\zeta)\|_{1,p,\alpha} + \|\bar{\partial}_J u_{(r)}\| \tag{174}$$

for some constant $C > 0$. For any small $\zeta \in kerD\mathcal{S}_b, exp_{u_i}\zeta$ converges to a periodic orbit as $|s_i| \to \infty$. It follows that $S_{exp_u \zeta}$ converges to zero exponentially. Since $(\kappa, h_{(r)})$ satisfies (171) with $\kappa|_{|s_1| \ge \frac{lr}{2kT}} = 0$, by elliptic estimate we conclude that in the part $\frac{lr}{2kT} \le |s_i| \le \frac{3lr}{2kT}$

$$|S_{exp_{u_{(r)}}h_{(r)}}| \le C_0 e^{-\mathfrak{c}\frac{lr}{2kT}}, \quad |S_{u_{(r)}}| \le C_0 e^{-\mathfrak{c}\frac{lr}{2kT}} \tag{175}$$

for r big enough. Moreover, we may choose r very large and $|(\kappa,\zeta)|$ very small such that

$$\|Q_{b_{(r)}}\| \left\| P_{exp_{u_{(r)}}h_{(r)}}\left(D\mathcal{S}_{exp_{u_{(r)}}h_{(r)}}(\Psi, G)\right) - D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}\left(\kappa_r, h_{(r)}\right)\right)\right\|_{1,p,\alpha,r}$$

$$\le \epsilon \left\| \frac{\partial}{\partial r}\left(\kappa_r, h_{(r)}\right)\right\|_{1,p,\alpha,r} + Ce^{-(\mathfrak{c}-\alpha)\frac{lr}{2kT}}, \tag{176}$$

where $\epsilon > 0$ is a constant such that $4\epsilon\mathcal{C}_3 \le 1$. Note that

$$\frac{\partial}{\partial r}(\kappa_r, h_{(r)}) = \frac{\partial}{\partial r}I_r(\kappa,\zeta) + \frac{\partial}{\partial r}\left(Q_{b_{(r)}}\right) \circ f_{(r)}(I_r(\kappa,\zeta)) + \left(Q_{b_{(r)}}\frac{\partial}{\partial r} \circ f_{(r)}(I_r(\kappa,\zeta))\right). \tag{177}$$

Then

$$D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}(\kappa_r, h_{(r)})\right) = D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}I_r(\kappa,\zeta)\right) + D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}(Q_{b_{(r)}})f_{(r)}(I_r(\kappa,\zeta))\right) + \frac{\partial}{\partial r}(f_{(r)}(I_r(\kappa,\zeta))).$$

It follows together with (174), (176) that

$$\begin{aligned} &\left\|\frac{\partial}{\partial r}(f_{(r)}(I_r(\kappa,\zeta)))\right\|_{p,\alpha,r} \\ \leq\ & \epsilon\left\|\frac{\partial}{\partial r}(\kappa_r, h_{(r)})\right\|_{1,p,\alpha,r} + \left\|D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}I_r(\kappa,\zeta)\right)\right\|_{p,\alpha,r} + \|D\mathcal{S}_{b_{(r)}} \circ \frac{\partial}{\partial r}(Q_{b_{(r)}}) \circ f_{(r)}(I_r(\kappa,\zeta))\|_{p,\alpha,r}. \end{aligned} \tag{178}$$

For any (κ_1, h_1), we have

$$\frac{\partial}{\partial r}\left(D\mathcal{S}_{b_{(r)}}\right)(\kappa_1, h_1) = \frac{\partial}{\partial r}\left(D_{u_{(r)}}h_1 + \kappa_1\right) = \frac{\partial S_{u_{(r)}}}{\partial r}h_1 = \left(0, \frac{\partial S_{u_{(r)}}}{\partial r}\right)\cdot(\kappa_1, h_1).$$

Then

$$\frac{\partial}{\partial r}\left(D\mathcal{S}_{b_{(r)}}\right) = \left(0, \frac{\partial S_{u_{(r)}}}{\partial r}\right) \tag{179}$$

As $I_r(\kappa,\zeta) \in ker D\mathcal{S}_{b_{(r)}}$ we have

$$D\mathcal{S}_{b_{(r)}} \circ I_r(\kappa,\zeta) = 0.$$

Then

$$D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}I_r(\kappa,\zeta)\right) + \left(0, \frac{\partial S_{u_{(r)}}}{\partial r}\right)I_r(\kappa,\zeta) = 0.$$

Since $\frac{\partial S_{u_{(r)}}}{\partial r}$ supports in the part $\frac{lr}{2kT} \leq |s_i| \leq \frac{3lr}{2kT}$, by the exponential decay of S_{u_i} we get

$$\left\|D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}I_r(\kappa,\zeta)\right)\right\|_{p,\alpha,r} \leq C_1 e^{-(\mathfrak{c}-\alpha)\frac{lr}{2kT}}. \tag{180}$$

Taking the differentiation of the equality

$$D\mathcal{S}_{b_{(r)}} \circ Q_{b_{(r)}} = I$$

we get

$$D\mathcal{S}_{b_{(r)}}\left(\frac{\partial}{\partial r}(Q_{b_{(r)}})\right) = -\left(0, \frac{\partial S_{u_{(r)}}}{\partial r}\right)\cdot Q_{b_{(r)}}.$$

Together with (174) and (175) we get

$$\|Q_{b_{(r)}}\|\left\|D\mathcal{S}_{b_{(r)}}\frac{\partial}{\partial r}(Q_{b_{(r)}})f_{(r)}(I_r(\kappa,\zeta))\right\|_{1,p,\alpha,r} \leq C_2 e^{-(\mathfrak{c}-\alpha)\frac{lr}{2kT}} \tag{181}$$

and

$$\left\|\frac{\partial}{\partial r}(f_{(r)}(I_r(\kappa,\zeta)))\right\|_{p,\alpha,r} \leq C_2 e^{-(\mathfrak{c}-\alpha)\frac{lr}{8kT}} + \epsilon\left\|\frac{\partial}{\partial r}\left(\kappa_r, h_{(r)}\right)\right\|_{1,p,\alpha,r}. \tag{182}$$

By (170) we have

$$\left\|\frac{\partial}{\partial r}\left(\kappa_r, h_{(r)}\right)\right\|_{1,p,\alpha,r} \leq \left(\left\|\frac{\partial}{\partial r}I_r(\kappa,\zeta)\right\|_{1,p,\alpha,r} + \left\|\frac{\partial}{\partial r}\left(Q_{b_{(r)}} \circ f_{(r)}(I_r(\kappa,\zeta))\right)\right\|_{1,p,\alpha,r}\right) \tag{183}$$

Together with (143) we have

$$\begin{aligned}
&\left\|\frac{\partial}{\partial r}\left(\kappa_r, h_{(r)}\right)\right\|_{1,p,\alpha,r} \\
&\leq \mathcal{C}_3\left(e^{-\alpha\frac{lr}{8kT}} + \left\|\frac{\partial}{\partial r} I_r(\kappa,\zeta)\right\| + \left\|\frac{\partial}{\partial r}\left(f_{(r)}(I_r(\kappa,\zeta))\right)\right\|_{1,p,\alpha,r} + \|f_r \circ I_r(\kappa,\zeta)\mid_{\frac{lr}{4kT}\leq|s_i|\leq\frac{7lr}{4kT}}\|_{p,\alpha,r}\right) \\
&\leq C_4 e^{-\alpha\frac{lr}{8kT}} + \mathcal{C}_3\epsilon\left\|\frac{\partial}{\partial r}\left(\kappa_r, h_{(r)}\right)\right\|_{1,p,\alpha,r}.
\end{aligned} \tag{184}$$

where we used (153), (157) and (182) in last inequality. Then the Lemma follows from $4\mathcal{C}_3\epsilon < 1$ and (184). □

10 Contact invariants and Open string invariants

Proposition 10.1. *U_I has the property*

(1). *Each strata of U_I is a smooth manifold.*

(2). *If $U_{I,D'} \subset U_{I,D}$ is a lower stratum, it is a submanifold of codimension at least 2.*

Proof. The proof of **(1)** for the top strata is standard, we omit it here. The proof for the lower strata will be given in our next paper [20]. It is well-known that an interior node corresponds to codimension 2 stratum. It suffices to consider those nodal corresponding to periodic orbits. Note that:

(1). we mod the freedoms of choosing the origin in $\mathbb{R}$ and the origin of the periodic orbits;

(2). we choose the Li-Ruan's compactification in [18], that is, we firstly let the Riemann surfaces degenerate in Delingne-Mumford space and then let M^+ degenerate compatibly. At any node, the Riemann surface degenerates independently with two parameters, which compatible with those freedoms of choosing the origins (see section §5 for degeneration and section §6 for gluing);

Then both blowups at interior and at infinity lead boundaries of codimension 2 or more. □

From section §7.2 we have a finite dimensional virtual orbifold system $\{U_I, E_I, \sigma_I\}$ and a finite dimensional virtual orbifold $\{U_I\}$ indexed by a partially ordered set ($\mathcal{I} = 2^{\{1,2,\cdots,N\}}, \subset$). By the same argument of Theorem 5.16 we conclude that $\{U_{I,\epsilon}(M^+)\}$ is a compact virtual manifold. As we have mod the S^1-action on periodic orbit, and S^1-action on the puncture points on Riemann surfaces, by the same method of [23] we can show that $\{U_{I,\epsilon}(M^+)\}$ is oriented (see [20]).

Let $\Lambda = \{\Lambda_I\}$ be a partition of unity and $\{\Theta_I\}$ be a virtual Euler form of $\{\mathbb{E}_I\}$ such that $\Lambda_I\Theta_I$ is compactly supported in $U_{I,\epsilon}$.

Recall that we have two natural maps

$$e_i : U_{I,\epsilon} \longrightarrow M^+$$

$$(u; \Sigma, \mathbf{y}, \mathbf{p}) \longrightarrow u(y_i)$$

for $i \leq m$ defined by evaluating at marked points and

$$e_j : U_{I,\epsilon} \longrightarrow Z_{\mathfrak{e}_j}$$

$$(u; \Sigma, \mathbf{y}, \mathbf{p}) \longrightarrow u(p_j)$$

for $j > m$ defined by projecting to its periodic orbit. The contact invariant can be defined as

$$\Psi^{(C)}_{(A,g,m+\nu,\mathbf{k},\mathfrak{e})}(\alpha_1, ..., \alpha_m; \beta_{m+1}, ..., \beta_{m+\nu}) = \sum_I \int_{U_{I,\epsilon}} \prod_i e_i^*\alpha_i \wedge \prod_j e_j^*\beta_j \wedge \Lambda_I\Theta_I. \tag{185}$$

for $\alpha_i \in H^*(M^+, \mathbb{R})$ and $\beta_j \in H^*(Z_{\mathfrak{e}_j}, \mathbb{R})$ represented by differential form. Clearly, $\Psi^C = 0$ if $\sum \deg(\alpha_i) + \sum \deg(\beta_i) \neq Ind^C$.

Similarly, the open string invariant can be defined as

$$\Psi^{(L)}_{(A,g,m+\nu,\vec{\mu})}(\alpha_1, ..., \alpha_m) = \sum_I \int_{U_{I,\epsilon}} \prod_i e_i^*\alpha_i \wedge \Lambda_I\Theta_I. \tag{186}$$

for $\alpha_i \in H^*(M^+, \mathbb{R})$. Clearly, $\Psi^{(L)} = 0$ if $\sum \deg(\alpha_i) \neq Ind^L$.

It is proved that these integrals are independent of the choices of Θ_I and the choices of the regularization (see [8]). We must prove the convergence of the integrals (185) and (186) near each lower strata.

Let $\alpha \in H^*(M^+, \mathbb{R})$ and $\beta \in H^*(Z, \mathbb{R})$ represented by differential form. We may write

$$\prod_i e_i^*\alpha_i \wedge \prod_j e_j^*\beta_j \wedge \Lambda_I\Theta_I = y dr \wedge d\tau \wedge dj \wedge d\zeta,$$

where $d\zeta$ and dj denote the volume forms of $ker D\mathcal{S}_b$ and the space of complex structures respectively and $y = y(r, \tau, j, \zeta)$ is a function. Then (169) implies that $|y| \leq C_1 e^{-C_2 r}$ for some constants $C_1 > 0,\ C_2 > 0$. Then the convergence of the integral (185) follows. Similarly, we can prove the convergence of the integral (186).

Obviously, both $\Psi^{(C)}$ and $\Psi^{(L)}$ are generalizations of the relative GW invariants.

One can easily show that

Theorem 10.2. *(i).* $\Psi^{(C)}_{(A,g,m+\nu,\mathbf{k},\mathfrak{e})}(\alpha_1, ..., \alpha_m; \beta_{m+1}, ..., \beta_{m+\nu})$ *is well-defined, multi-linear and skew symmetric.*

(ii). $\Psi^{(C)}_{(A,g,m+\nu,\mathbf{k},\mathfrak{e})}(\alpha_1, ..., \alpha_m; \beta_{m+1}, ..., \beta_{m+\nu})$ *is independent of the choice of forms* α_i, β_j *representing the cohomology classes* $[\beta_j], [\alpha_i]$*, and the choice of virtual neighborhoods.*

(iii). $\Psi^{(C)}_{(A,g,m+\nu,\mathbf{k},\mathfrak{e})}(\alpha_1, ..., \alpha_m; \beta_{m+1}, ..., \beta_{m+\nu})$ *is independent of the choice of* $\widetilde{J}$ *and* J *over* M_0^+.

Theorem 10.3. *(i).* $\Psi^{(L)}_{(A,g,m+\nu,\vec{\mu})}(\alpha_1, ..., \alpha_m)$ *is well-defined, multi-linear and skew symmetric.*

(ii). $\Psi^{(L)}_{(A,g,m+\nu,\vec{\mu})}(\alpha_1, ..., \alpha_m)$ *is independent of the choice of forms* α_i *representing the cohomology classes* $[\alpha_i]$*, and the choice of virtual neighborhoods.*

(iii). $\Psi^{(L)}_{(A,g,m+\nu,\vec{\mu})}(\alpha_1, ..., \alpha_m)$ *is independent of the choice of* $\widetilde{J}$ *and* J *over* M_0^+.

References

[1] M. Aganagic, A. Klemm, C. Vafa, Disk instantons, mirror symmetry and the duality web, Z. Naturforsch. A 57 (2002) 1C28

[2] M. Aganagic, C. Vafa, Mirror Symmetry, D-Branes and Counting Holomorphic Discs. arxiv.org:hep-th/0012041v1, 2000

[3] A. Adem, J. Leida and Y. Ruan, Orbifolds and string topology, Camridge Tracts in Mathematics, 171. Cambridge University Press, Cambridge, 2007.

[4] K. Behrend, *Gromov-Witten invariants in algebraic geometry.* Invent. Math. 127(3), 601-617, 1997.

[5] F. Bourgeois, A Morse-Bott approach to contact homology, Ph.D. thesis, Stanford University, 2002

[6] Bourgeois, F., Eliashberg, Y., Hofer, H., Wysocki, K., Zehnder, E., *Compactness results in symplectic field theory,* Geom. Topol. **7** (2003), 799-888.

[7] B. Chen, A. Li and B. Wang, Virtual neighborhood technique for pseudo-holomorphic spheres, arXiv:1306.32

[8] B. Chen and G. Tian, Virtual manifolds and Localization, Acta Math. Sinica, 2010(26),1-24.

[9] S. K, Donaldson, Floer homology groups in Yang-Mills theory. With the assistance of M. Furuta and D. Kotschick. Cambridge Tracts in Mathematics, 147. Cambridge University Press, Cambridge, 2002. viii+236 pp.

[10] K. Fukaya and K. Ono, Arnold conjecture and Gromov-Witten invariants, Topology 38,1999,933-1048.

[11] B. Fang and C.-C. M. Liu, Open Gromov-Witten invariants of toric Calabi-Yau 3-folds, Comm. Math. Phys. 323 (2013), no. 1, 285-328.

[12] M. Gromov, Pseudo holomorphic curves in symplectic manifolds, Invent. math., 82 (1985), 307-347.

[13] H. Hofer, Pseudoholomorphic curves in symplectizations with applications to the weinstein conjecture in dimension three, Invent. Math. 114(1993), 515-563.

[14] H. Hofer, K. Wysocki, E. Zehnder,Properties of pseudoholomorphic curves in symplectisations. I. Asymptotics. Ann. Inst. H. Poincar Anal. Non Linaire 13 (1996), no. 3, 337-379.

[15] H. Hofer, K. Wysocki, E. Zehnder, Properties of pseudo-holomorphic curves in symplectisations. II. Embedding controls and algebraic invariants. Geom. Funct. Anal. 5 (1995), no. 2, 270-328.

[16] S. Katz, C.-C.M. Liu, Enumerative geometry of stable maps with Lagrangian boundary conditions and multiple covers of the disc. The interaction of finite-type and Gromov-Witten invariants (BIRS 2003), 1-47, Geom. Topol. Monogr., 8, Geom. Topol. Publ., Coventry, 2006.

[17] J. Li, C.-C.M. Liu,K.Liu and J.Zhou, A mathematical theory of the topological vertex, Geom. Topol. 13(2009), no.1,527-621.

[18] A-M. Li and Y. Ruan, Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds, Invent. Math. 145, 151-218(2001)

[19] A-M. Li and Li Sheng, Exponential decay of J-holomorphic maps with finite energy, in preparation.

[20] A-M. Li and Li Sheng, Relative invariants, contact geometry and open string invariants-2, in preparation.

[21] J. Li and G. Tian, Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties. J. Amer. Math. Soc., 11(1), 119-174,1998.

[22] G. Liu and G. Tian, Floer homology and Arnold conjecture. J. Differential Geom. 49 (1998), no. 1, 1-74.

[23] D. McDuff and D. Salamon, J-holomorphic curves and quantum cohomology, University Lec. Series, vol. 6, AMS.

[24] D. McDuff and D. Salamon, Introduction to symplectic topology, Oxford Press.

[25] T. Parker and J. Woflson, A compactness theorem for Gromov's moduli space, J. Geom. Analysis, 3 (1993), 63-98.

[26] Y. Ruan, Surgery, quantum cohomology and birational geometry, Northern California Symplectic Geometry Seminar, 183-198, Amer. Math. Soc. Transl. Ser. 2, 196.

[27] Y. Ruan, Virtual neighborhoods and pseudo-holomorphic curves, Turkish Jour. of Math. 1(1999), 161-231.

[28] Y. Ruan, G. Tian, A mathematical theory of quantum cohomology. J. Diff Geom. 42(1995),259-367.

[29] B. Siebert, Gromov-Witten invariants for general symplectic manifolds, arXiv:9608005

[30] S. Lang, Real analysis, second edition,S Addison-Wesley Publishing Company, Advanced Book Program, Reading, MA, 1983. xv+533 pp.

[31] H. Ooguri, C. Vafa, Knot invariants and topological strings. Nuclear Phys. B 577 (2000), no. 3, 419-438.

[32] Y.-G. Oh, R. Wang, Analysis of contact Cauchy-Riemann maps II: canonical neighborhoods and exponential convergence for the Morse-Bott case, arXiv:1311.6196